



REC'D 2 7 AUG 2003

WIPO

PCT

Patent Office Canberra

I, LEANNE MYNOTT, MANAGER EXAMINATION SUPPORT AND SALES hereby certify that annexed is a true copy of the Provisional specification in connection with Application No. 2002951579 for a patent by MELBOURNE HEALTH as filed on 22 August 2002.



WITNESS my hand this Eighteenth day of August 2003

LEANNE MYNOTT

MANAGER EXAMINATION SUPPORT

AND SALES

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

Regulation 3.2

Melbourne Health

# AUSTRALIA Patents Act 1990

## PROVISIONAL SPECIFICATION

for the invention entitled:

"Diagnostic and therapeutic agents"

The invention is described in the following statement:

### DIAGNOSTIC AND THERAPEUTIC AGENTS

#### FIELD OF THE INVENTION

The present invention relates generally to diagnostic and therapeutic agents. More particularly, the present invention provides mammalian transcription factors which function in the modulation of expression of genetic sequences. The present invention further provides nucleic acid molecules encoding the transcription factors as well as nucleic acid and/or proteinaceous molecules with which the transcription factors interact.

The transcription factors of the present invention or molecules interacting with same may be used *inter alia* in the generation of a range of diagnostic and therapeutic agents for a range of conditions.

#### BACKGROUND OF THE INVENTION

Reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common

general knowledge in any country.

20 Bibliographic details of references provided in the subject specification are listed at the end of the specification.

of the specification.

15

The increasing sophistication of recombinant DNA techniques has provided significant progress in understanding the mechanisms involved in regulating eukaryotic gene expression. This is greatly facilitating research and development in the plant, agricultural, medical and veterinary industries. Transcription factors are an important component in the control of gene expression. However, despite their importance, mammalian transcription factors have not been well investigated for their diagnostic and therapeutic potential.

30 RNA polymerases in eukaryotic cells cannot initiate transcription alone; before transcription can begin, they require interaction between transcription factors and the

20

5%

promoter. These factors assemble at the promoter and, via a series of steps, facilitate both the binding of RNA polymerase II to the promoter and its subsequent phosphorylation and release to initiate transcription.

In addition to these general transcription factors, many thousands of transcription activators and/or negative regulators (inhibitors) exist, which control the process of initiation of gene transcription from great distances along the DNA. These factors influence the timing and extent of transcription of a particular gene. Indeed, they control whether and to what extent a particular gene is transcribed in a cell of a particular tissue type. Although most gene regulators identified to date have been found to be proteins, some transcription factors may also be RNA molecules.

In *Drosophila*, the transcription factor known as "Grainyhead" regulates key developmental process in the embryo and is encoded by the gene grainyhead. During development, Grainyhead is initially involved in dorsal/ventral and terminal patterning of the newly fertilized embryo through the formation of multi-protein complexes that repress transcription from the decapentaplegic, tailless and zerknuellt genes (Huang et al., Genes Dev. 9: 3177-3189, 1995; Liaw et al., Genes Dev. 9: 3163-3176, 1995). Later, grainyhead is predominantly expressed in the embryonic central nervous system in cuticle-producing tissues, where it binds to promoters and influences transcription from other developmentally regulated genes including engrailed, fushi tarazu and Ultrabithorax (Bray et al., Genes Dev. 3: 1130-1145, 1989; Dynlacht et al., Genes Dev. 3: 1677-1688, 1989; Biggin and Tjian, Cell 53: 699-711, 1988; Soeller et al., Genes Dev. 2: 68-81, 1988; Dynlacht et al., Cell 56: 563-576, 1991; Attardi and Tjian, Genes Dev. 7: 1341-1353, 1993; Uv et al., Mol. Cell Biol. 14: 4020-4031, 1994).

The importance of grainyhead in Drosophila development is emphasised by the embryonic lethal phenotype observed in flies carrying mutations in this gene. The embryos have flimsy cuticles, grainy and discontinuous head skeletons and patchy tracheal tubes (Bray and Kafatos, Genes Dev. 5: 1672-1683, 1991). A neuroblast-specific isoform of the protein, arising from alternate splicing, has also been identified. A mutation that abolishes

this isoform is pupal- and adult- lethal, and flies demonstrate uncoordinated movements (Uv et al., Mol. Cell Biol. 17: 6727-6735, 1997).

Mammalian homologs of grainyhead have previously been proposed, including three genes designated CP2, LBP-1a and LBP-9. Studies have implicated them in a wide variety of cellular and developmental events including T cell proliferation, globin gene expression and steroid biosynthesis (Sueyoshi et al., Mol. Cell Biol. 15: 4158-4166, 1995; Jane et al., EMBO J. 14: 97-105, 1995; Volker et al., Genes Development 11: 1435-1446, 1997; Zhou et al., Mol. Cell Biol. 20: 7662-7672, 2000). However, in situ analyses of both CP2 and 10 LBP-1a reveal ubiquitous expression of both genes, unlike the highly restricted pattern observed with grainyhead in Drosophila (Bray et al., 1989, supra; Dynlacht et al., 1989, supra; Bray and Kafatos, 1991, supra; Ramamurthy et al., J. Biol. Chem. 276: 7836-7842, 2001). It is concluded, therefore, that these genes are not close homologs of grainyhead.

15 There is a need to identify other mammalian transcription factors and in particular close mammalian homologs of Grainyhead and to use these to develop a range of diagnostic and therapeutic agents.

25

#### SUMMARY OF THE INVENTION

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the 5 inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

Nucleotide and amino acid sequences are referred to by a sequence identifier number (SEQ ID NO:). The SEQ ID NOs: correspond numerically to the sequence identifiers <400>1 10 '(SEQ ID NO:1), <400>2 (SEQ ID NO:2), etc. A sequence listing is provided at the end of the specification. A summary of the SEQ ID NOs is provided in Table 1.

Genetic sequences were studied exhibited homology at the nucleotide and/or amino acid level to a Drosophila gene, the product of which is involved in body patterning where a fine balance between activation and inhibition of gene expression is critical to the correct development of cells and tissues into functional organisms. A large number of different families of transcription factors play a critical role in ensuring that this balance is maintained during embryological development. One such transcription factor, cloned from Drosophila and well-characterized, is Grainyhead (hereinafter referred to by its 20 abbreviation, GRH). GRH is encoded by the gene grainyhead (grh). The inventors observed that the identity of previously published putative grh mammalian homologs showed much more ubiquitous expression compared with the highly restricted pattern exhibited by Drosophila grh. Furthermore, sequence similarity between the proposed mammalian homologs and the Drosophila grh sequence was relatively low. In accordance with the present invention, true grh homologs were identified and derived from mammalian tissue such as human and mouse tissue.

Accordingly, one aspect of the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a mammalian homolog of Drosophila GRH. A mammalian homolog of GRH is referred to herein as M-GRH. The corresponding gene is referred to as M-grh. A M-grh is

15

deemed a homolog of *Drosophila grh* (*D-grh*). If it comprises a nucleotide sequence having 60% or greater similarity to the nucleotide sequence of *D-grh* after optimal alignment. Likewise, a M-GRH is so defined if it comprises an amino acid sequence having 60% or greater similarity to the amino acid sequence of *Drosophila* GDH (*D-GRH*). There are four isoforms of *Drosophila grh* designated *D-grh* P1, *D-grh* P2, *D-grh* P3 and *D-grh* P4. The nucleotide sequence encoding *D-grh* is set forth in SEQ ID NO:17 and SEQ ID NO:34, SEQ ID NO:36 and SEQ ID NO:38, respectively. Mammalian sequences encompassed by the present invention include those derived from tissues of mouse and human including, for example, mouse embryo, human fetal brain and placenta, and mouse and human kidney. Reference herein to *Drosophila grp* includes any or all of its isoforms P1-P4.

The mammalian sequences identified by the present inventors show higher percentages of similarity to the *D-grh* sequence than the already identified mammalian sequences designated *CP2*, *LBP-1a* and *LBP-9*. In accordance with the present invention, it is proposed that the M-grh homologs disclosed are "true" grh homologs relative to *CP2*, *LBP-1a* and *LBP-9*. As a result of the analysis herein described, it is shown that the earlier sequences align phylogenetically with another distinct *Drosophila* factor, designated *Drosophila CP2*. A new family of transcription factors, highly conserved from *Drosophila* to human and having distinct tissue-specificity profiles, is now described in accordance with the present invention.

The true M-grh homologs of the present invention include mammalian grainyhead (gene: mgr; expression product: MGR), brother of mgr (gene: bom; expression product: BOM) and sister of mgr (gene som: protein: SOM). MGR has multiple isoforms including MGR p49 and MGR p70 in humans and MGR p61 in mice. A summary of the SEQ ID NOs for the M-grh and M-GRH molecules of the present invention are shown in Table 2. The sequences are provided in the Sequence Listing.

30 The present invention provides, therefore, expression products of the M-grh genes, mgr, bom and som as well as derivatives and homologs thereof. This aspect of the present

invention does not extend to CP2, LBP-1a or LBP-9.

Accordingly, another aspect of the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding a polypeptide comprising a predicted amino acid sequence substantially as set forth in SEQ ID NO:2 (human MGR p49), SEQ ID NO:4 (human MGR p70), SEQ ID NO:6 (human BOM), SEQ ID NO:8 (human SOM), SEQ ID NO:10 (murine MGR p49), SEQ ID NO:12 (murine MGR p70), SEQ ID NO:14 (murine BOM) or SEQ ID NO:16 (murine SOM) or an amino acid sequence having at least about 60% similarity to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16 after optimal alignment.

The preferred nucleic acid molecules comprise sequences of nucleotides substantially as set forth in SEQ ID NO:1 (human mgr p49), SEQ ID NO:3 (human mgr p70), SEQ·ID NO:5 (human bom), SEQ ID NO:7 (human som), SEQ ID NO:9 (murine mgr p61), SEQ ID NO:11 (murine mgr p70), SEQ ID NO:13 (murine bom) or SEQ ID NO:15 (murine som) or complementary forms thereof, or a nucleotide sequence having at least about 60% similarity to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 after optimal alignment or their complementary forms or a nucleotide sequence capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or complementary forms thereof under low stringency conditions. Again, this aspect of the present invention does not extend to nucleic acid molecules encoding CP2, LBP-1 and LBP-9.

25

20

The present invention further extends to recombinant forms of the M-GRH molecules. Preferred recombinant M-GRH molecules having amino acid sequences defined in parenthesis include human MGR p49 (SEQ ID NO:2), human MGR p70 (SEQ ID NO:4), human BOM (SEQ ID NO:6), human SOM (SEQ ID NO:8), murine MGR p61 (SEQ ID NO:10), murine MGR p70 (SEQ ID NO:12), murine BOM (SEQ ID NO:14) and murine SOM (SEQ ID NO:16).

25

30

Reference to "M-GRH" molecules include derivatives, homologs and analogs thereof.

The mammalian transcription factors of the present invention are proposed to be involved in the regulation of expression of a range of genes such as but not limited to developmentally regulated genes involved in determining patterning. Some of the genes regulated encode critical products, the absence or malfunctioning of which, is proposed to lead to unwanted phenotypes and/or predispositions to certain medical conditions. That is, the presence of a mutation in and/or malfunction of a M-grh including over or under expression of the transcription factors of the present invention are proposed to cause incorrect regulation of one or more of these genes thereby leading to an inappropriate phenotype. The ability to detect mutations in the nucleotide sequences encoding the M-grh homologs permits the detection of a range of abnormalities or a predisposition for development of abnormalities. Furthermore, as many of the genes will be developmentally regulated genes, identification of the transcription factors permits identification of unknown developmentally regulated genes.

Accordingly, another aspect of the present invention contemplates a method for detecting a variation in a polynucleotide sequence encoding a M-GRH transcription factor.

Furthermore, the isolated nucleic acid molecules of the present invention may be able to be used to correct such an abnormality in a subject in need thereof or at risk of developing an abnormality. The nucleic acid molecules of the present invention may be comprised, therefore, within a suitable vector for delivery of all or part of the sequence to a recipient cell or tissue. The nucleic acid molecule or part thereof could also be administered directly for transient expression. The present invention provides, therefore, the potential for both a diagnostic and a therapeutic capability.

Accordingly, a further aspect of the present invention contemplates a genetic construct comprising a nucleotide sequence selected from SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15 or a

nucleotide sequence having at least 60% similarity to one or more of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 after optimal alignment or a nucleotide sequence capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or a complementary form thereof under low stringency conditions.

In a related embodiment, the present invention provides a genetic construct comprising a promoter or functional equivalent thereof operably linked to a nucleotide sequence of the invention.

Genes are represented herein in lower case italics. Expression products (e.g. proteins or RNA) are represented in upper case, non-itallic letters. A summary of the genes and their expression products is provided in Table 1.

TABLE 1
Abbreviations

GENE	EXPRESSION PRODUCT	
grainyhead (grh)	Grainyhead (GRH)	
mammalian grainyhead homologs (M-grh)	mammalian grainyhead homologs (M-GRH)	
mammalian grainyhead (mgr)	mammalian Grainyhead (MGR)	
brother of mammalian grainyhead (bom)	brother of mammalian grainyhead (BOM)	
sister of mammalian grainyhead (som)	sister of mammalian grainyhead (SOM)	

15

10

A summary of sequence identifiers used throughout the specification is Table 2.

TABLE 2
Summary of sequence identifiers

5

SEQUENCE	NAME	DESCRIPTION
ID NO.		
1	human <i>mgr</i> p49	Nucleotide sequence encoding mammalian grainyhead derived from human fetal brain
2	human MGR p49	Predicted amino acid sequence corresponding to SEQ ID NO:1
3	human <i>mgr</i> p70	Nucleotide sequence encoding mammalian grainyhead being an isoform of SEQ ID NO:1, derived from human kidney
4	human MGR p70	Predicted amino acid sequence corresponding to SEQ ID NO:3
5	human bom	Nucleotide sequence encoding mammalian grainyhead derived from human placenta
6	human BOM	Predicted amino acid sequence corresponding to SEQ ID NO:5
7	human som	Nucleotide sequence encoding mammalian grainyhead
8	human SOM	Predicted amino acid sequence corresponding to SEQ ID NO:7
9	murine <i>mgr</i> p61	Nucleotide sequence encoding mammalian grainyhead derived from 17.5 day murine embryo
10	murine MGR p61	Predicted amino acid sequence corresponding to SEQ ID NO:9
11	murine <i>mgr</i> p70	Nucleotide sequence encoding mammalian grainyhead being an isoform of SEQ ID NO:9, derived from murine kidney
12	murine MGR p70	Predicted amino acid sequence corresponding to SEQ ID NO:11
13	murine bom	Nucleotide sequence encoding mammalian grainyhead derived from a murine embryonic carcinoma cell line (p19)
14	murine BOM	Predicted amino acid sequence corresponding to SEQ ID NO:13
15	murine som	Nucleotide sequence encoding mammalian grainyhead
16	murine SOM	Predicted amino acid sequence corresponding to SEQ ID NO:15
17	grh-P1	Nucleotide sequence encoding the Drosophila

SEQUENCE ID NO.	NAME	DESCRIPTION
		transcription factor designated Grainyhead (grh)
18	GRH-P1	Amino acid sequence corresponding to SEQ ID NO:18
19-20	human p49 mgr	primers
21-22	human p70 <i>mgr</i>	primers
23-24	human <i>bom</i>	primers
25-26	murine p70 <i>mgr</i>	primers
27-28	murine p61 mgr	primers
29-30	murine bom	primers
31-32	human S14	primers
33	Drosophila dopa decarboxylase	promoter
34	Drosophila PCNA	promoter
35	human Engrailed-1	promoter
36	grh-P2	Nucleotide sequence encoding the <i>Drosophila</i> transcription factor designated <i>Grainyhead</i> (grh) isoform P2
37	GRH-P2	Amino acid sequence corresponding to SEQ ID NO:36
38	grh-P3	Nucleotide sequence encoding the Drosophila transcription factor designated Grainyhead (grh) isoform P3
39	GRH-P3	Amino acid sequence corresponding to SEQ ID NO:38
40	grh-P4	Nucleotide sequence encoding the Drosophila transcription factor designated Grainyhead (grh) isoform P4
41	GRH-P4	Amino acid sequence corresponding to SEQ ID NO:40

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a representation showing that mgr genomic locus encodes two distinct isoforms. (A) Alignment of the predicted NH2-terminal amino acid sequence of the p70 5 isoform of MGR and BOM. Amino acid identity is denoted by shared upper case letters and similarity by the (+) symbol. The first amino acids shared between p61 MGR and p70 MGR are given in bold. (B) Structure of the human and murine mgr genomic loci. Human genomic sequence was downloaded from the GenBank database (Accession Number AC010969) and aligned with cDNA sequences. Murine genomic clones were obtained from a 129 library and mapped by Southern analysis and PCR. Exons are denoted as E1-8 in human and E1-9 in murine. The two human MGR isoforms are denoted as p70 and p49 MGR and the two murine isoforms as p70 and p61 MGR. The scale of 1 kb is shown. (C) Identification of the murine p61 MGR promoter. Sequence was obtained from intron three from the MGR genomic locus and analyzed using the weight matrices of Bucher, J. Mol. Biol. 212: 563-578, 1990. The CAP site, TATA box and GC box are indicated. The cDNA start site is shown in arrows, the first ATG is given in bold and the splice site at the end of the first exon of p61 MGR is indicated.

Figure 2 is a photographic representation showing that p70 MGR binds to Drosophila gene regulatory sequences which bind grh. (A) p70 MGR binds to the Drosophila PCNA 20 promoter. Nuclear extract from the JEG-3 cell line was studied in an EMSA with a PCNA promoter probe in the presence and absence of anti-MGR specific antisera. Antisera 611 was raised against peptides common to the p70 and p49 MGR proteins in the dimerization domain and antisera 67 was raised against unique peptides in the NH2-terminal domain of p70 MGR. The migration of the MGR complex is shown in arrows. (B) p70 MGR binds to the Drosophila dopo decarboxylase promoter. Experimental conditions were as described for (A).

Figure 3 are representations showing that p70 MGR binds to and transactivates the human En-1 promoter. (A) Identification of a grh consensus DNA binding site in the human En-1 promoter. The consensus sequence for grh DNA binding compiled from an alignment of the Drosophila Ultrabithorax, Dopa decarboxylase and fushi tarazu promoters was compared with the sequence of the proximal human En-1 promoter and the Drosophila engrailed promoter. The closed bracket indicates the extend of the grainyhead binding site in the engrailed promoter as defined by DNAseI footprinting. (B) Human p70 MGR binds to the human En-1 promoter. Nuclear extract from the JEG-3 cell line was studied in an EMSA with a Ddc promoter probe in the presence of pre-immune sera (lane 1), anti-MGR specific antisera 67 (detailed in legend to Figure 2) (lane 2) or cold competitor DNA (lanes 3-5). A 50-fold excess of the Ddc probe was used in lane 3 and a 10- and 20-fold excess of a human En-1 promoter probe in lanes 4 and 5, respectively. The migration of the MGR/DNA complex is shown by arrows. (C) Human p70 MGR transactivates the En-1 promoter. COS cells were transiently transfected with the proximal En-1 promoter containing the MGR binding site linked to a mimimal y-globin promoter and a firefly luciferase reporter gene (solid columns), the miminal γ-globin promoter/luciferase reporter gene (open columns) and the TK promoter linked to the Renilla luciferase reporter gene (hatched columns) in the presence and absence of a p70 MGR expression vector (PCI-p70 MGR) as indicated. Transfection with the empty vector (pCI) served as the control. Luciferase levels were corrected for protein concentration and values were derived from two independent experiments performed in triplicate.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is predicated in part on the identification of mammalian homologs of the *Drosophila* transcription factor known as Grainyhead (GRH). GRH is encoded by the gene, grainyhead (grh) In *Drosophila*, mutations in this gene are associated with embryonic lethal phenotypes, indicating the importance of the gene for normal development and function. The mammalian homologs are proposed to be involved in the regulation of developmental and/or non-developmental genes. Identification and isolation of the mammalian homologs of grh (M-grh) enable the development of a range of diagnostic and therapeutic agents useful in the detection and treatment of genetic disorders.

The present invention provides, therefore, a family of mammalian-derived transcription factors, highly related from Drosophila to mammals. These transcription factors are more highly conserved than CP2, LBP-1a and LBP-9. The present invention does not extend to CP2, LBP-1 and LBP-9. Reference to a mammal in this context includes a human, livestock animal (e.g. sheep, cow, horse, pig, donkey, goat), laboratory test animal (e.g. mouse, rat, rabbit, guinea pig), companion animal (e.g. dog, cat) or captive wild animal. Most preferably, the animal is a human or murine species. Sources of the isolated nucleic acid molecules include a range of tissues, such as mouse embryo, human fetal brain and placenta, and mouse and human kidney. In view of the highly conserved nature of this family of M-grh nucleotide sequences, however, corresponding homologs from other tissues and from other mammalian species are intended to be included within the scope of the present invention. The term "homolog" as used herein, therefore, extends to encompass transcription factors from mammalian species encoded by nucleotide sequences which have substantial similarity to Drosophilia grh or a conserved region thereof. At the protein level, a homolog includes an amino acid sequence and/or tertiary structure having similarity to Drosophila GRH. In cases where the expression product of the M-grh is RNA, a homolog is defined by reference to the similar ribonucleotide sequence to that encoded by Drosophila grh.

30

10

M-ghd or M-GRH, i.e. a mammalian homolog of Drosophila grh or GRH is defined as

15

20

25

such by having a nucleotide or amino acid sequence which has 60% or greater similarity after optimal alignment to *Drosophila grh* or GRH.

Accordingly, one aspect of the present invention provides an isolated nucleic acid molecule comprising a sequence of nucleotides encoding or complementary to a sequence encoding a mammalian homolog of *Drosophila grh*.

Reference to a mammalian homolog of *Drosophila* GRH (i.e. a M-GRH) preferably includes the mammalian homolog of grainyhead (MGR), brother of MGR (BOM) and sister of MGR (SOM). These transcription factors are encoded by *mgr*, *bom* and *som*, respectively. Reference to "MGR", "BOM" and "SOM" or *mgr*, *bom* and *som* includes all mutants, derivatives, homologs and analogs thereof. The present invention further extends, however, to all novel mammalian homologs of *Drosophila grh* but does not encompass CP2, LBP-1a or LBP-9. The nucleotide sequences for *Drosophila grh* are set forth in SEQ ID NO:17, SEQ ID NO:34, SEQ ID NO:36 and SEQ ID NO:38, respectively. Consequently, a mammalian homolog is defined herein as comprising a nucleotide sequence having at least about 60% sequence similarity to SEQ ID NO:17 or SEQ ID NO:36 or SEQ ID NO:36 or SEQ ID NO:38 after optimal alignment and/or being capable of hybridizing to SEQ ID NO:17 or SEQ ID NO:34 or SEQ ID NO:36 or SEQ ID NO:38 or its complementary form under low stringency conditions.

Accordingly, another aspect of the present invention provides an isolated nucleic acid molecule encoding a mammalian transcription factor or a functional part thereof comprising a sequence of nucleotides having at least 60% similarity to SEQ ID NO:17 or SEQ ID NO:34 or SEQ ID NO:36 or SEQ ID NO:38 after optimal alignment and/or being capable of hybridizing to SEQ ID NO:17 or its complementary form under low stringency conditions.

In a preferred embodiment, the isolated nucleic acid molecule encodes a proteinaceous form of a transcription factor. Examples of such mammalian protein transcription factors include human MGR p49 (SEQ ID NO:2), human MGR p70 (SEQ ID NO:4), human

20

30

BOM (SEQ ID NO:6), human SOM (SEQ ID NO:7), murine MGR p61 (SEQ ID NO:10), murine MGR p70 (SEQ ID NO:12), murine BOM (SEQ ID NO:14) and murine SOM (SEQ ID NO:16).

Accordingly, another aspect of the present invention is directed to an isolated nucleic acid molecule comprising a sequence of nucleotides encoding a polypeptide having transcription factor activity and comprising an amino acid sequence substantially as set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 or SEQ ID NO:16 or an amino acid sequence having at least about 60% similarity to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 or SEQ ID NO:16 after optimal alignment wherein said polypeptide is a mammalian homolog of Drosophia GRH.

Such a polypeptide is referred to herein as a M-GRH.

Preferred percentage amino acid similarity levels include at least about 61% or at least about 62% or at least about 63% or at least about 64% or at least about 65% or at least about 66% or at least about 67% or at least about 68% or at least about 69% or at least about 70% or at least about 71% or at least about 72% or at least about 73% or at least about 74% or at least about 75% or at least about 76% or at least about 77% or at least about 78% or at least about 79% or at least about 80% or at least about 81% or at least about 82% or at least about 83% or at least about 84% or at least about 85% or at least about 86% or at least about 87% or at least about 88% or at least about 89% or at least about 90% or at least about 91% or at least about 92% or at least about 93% or at least about 94% or at least about 95% or at least about 96% or at least about 97% or at least 25 about 98% or at least about 99% similarity.

This aspect of the present invention includes derivatives of M-GRH molecules. Such derivatives include non-active fragments which encompass inter alia the binding domain as well as active isoforms.

15

20

A "derivative" of a polypeptide of the present invention also encompasses a portion or a part of a full-length parent polypeptide, which retains the transcription factor activity of the parent polypeptide. Such "biologically-active fragments" include deletion mutants and small peptides, for example, of at least 10, preferably at least 20 and more preferably at least 30 contiguous amino acids, which exhibit the requisite activity. Peptides of this type may be obtained through the application of standard recombinant nucleic acid techniques or synthesized using conventional liquid or solid phase synthesis techniques. For example, reference may be made to solution synthesis or solid phase synthesis as described, for example, in Chapter 9 entitled "Peptide Synthesis" by Atherton and Shephard which is included in a publication entitled "Synthetic Vaccines" edited by Nicholson and published by Blackwell Scientific Publications. Alternatively, peptides can be produced by digestion of an amino acid sequence of the invention with proteinases such as endoLys-C, endoArg-C, endoGlu-C and staphylococcus V8-protease. The digested fragments can be purified by, for example, high performance liquid chromatographic (HPLC) techniques. Any such fragment, irrespective of its means of generation, is to be understood as being encompassed by the term "derivative" as used herein.

In another embodiment, the present invention provides an isolated nucleic acid molecule encoding a mammalian transcription factor homolog of *Drosophila grh* (i.e. a M-GRH) and comprising a nucleotide sequence selected from SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 and SEQ ID NO:15 or a nucleotide sequence having at least about 60% similarity to any one of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 after optimal alignment or a nucleotide sequence capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:13 or SEQ ID NO:15 or a complementary form thereof under low stringency conditions.

Preferably, percentage nucleotide similarity levels include at least about 61% 61% or at least about 62% or at least about 63% or at least about 64% or at least about 65% or at least about 66% or at least about 67% or at least about 68% or at least about 69% or at

least about 70% or at least about 71% or at least about 72% or at least about 73% or at least about 74% or at least about 75% or at least about 76% or at least about 77% or at least about 78% or at least about 80% or at least about 81% or at least about 82% or at least about 83% or at least about 84% or at least about 85% or at least about 86% or at least about 87% or at least about 88% or at least about 89% or at least about 90% or at least about 91% or at least about 92% or at least about 93% or at least about 94% or at least about 95% or at least about 96% or at least about 97% or at least about 98% or at least about 99% similarity.

- 10 The term "similarity" as used herein includes exact identity between compared sequences at the nucleotide or amino acid level. Where there is non-identity at the nucleotide level, "similarity" includes differences between sequences which result in different amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. Where there is non-identity at the amino acid level, "similarity" includes amino acids that are nevertheless related to each other at the structural, functional, biochemical and/or conformational levels. In a particularly preferred embodiment, nucleotide and sequence comparisons are made at the level of identity rather than similarity.
- Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include "reference sequence", "comparison window", "sequence similarity", "sequence identity", "percentage of sequence similarity", "percentage of sequence identity", "substantially similar" and "substantial identity". A "reference sequence" is at least 12 but frequently 15 to 18 and often at least 25 or above, such as 30 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (i.e. only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window" refers to a conceptual segment of typically

12 contiguous residues that is compared to a reference sequence. The comparison window may comprise additions or deletions (i.e. gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, WI, USA) or by inspection and the best alignment (i.e. resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul *et al.* (*Nucl. Acids. Res.* 25: 3389, 1997). A detailed discussion of sequence analysis can be found in Unit 19.3 of Ausubel *et al.* (In: Current Protocols in Molecular Biology, John Wiley & Sons Inc. 1994-1998).

The terms "sequence similarity" and "sequence identity" as used herein refers to the extent that sequences are identical or functionally or structurally similar on a nucleotide-bynucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a "percentage of sequence identity", for example, is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g. A, T, C, G, I) or the identical amino acid residue (e.g. Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. For the purposes of the present invention, "sequence identity" will be understood to mean the "match percentage" calculated by the DNASIS computer program (Version 2.5 for windows; available from Hitachi Software engineering Co., Ltd., South San Francisco, California, USA) using standard defaults as used in the reference manual accompanying the software. Similar comments apply in relation to sequence similarity.

30

15

20

25

The present invention provides, therefore, an isolated nucleic acid molecule comprising a

sequence of nucleotides selected from SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 and SEQ ID NO:15 or a complementary form thereof. Such nucleic acid molecules encode mammalian homologs of *Drosophila grh*. These mammalian homologs are proposed herein to be transcription factors.

The present invention extends to variants of the nucleic acid molecules. A variant is a molecule having less than 100% sequence identity to a M-grh. Generally, a variant will still hybridize to a M-grh sequence under low stringency conditions.

10

5

The term "variant" refers, therefore, to nucleotide sequences displaying substantial sequence identity with a reference nucleotide sequences or polynucleotides that hybridize with a reference sequence under stringency conditions that are defined hereinafter. The terms "nucleotide sequence", "polynucleotide" and "nucleic acid molecule" may be used herein interchangeably and encompass polynucleotides in which one or more nucleotides have been added or deleted, or replaced with different nucleotides. In this regard, it is well understood in the art that certain alterations inclusive of mutations, additions, deletions and substitutions can be made to a reference nucleotide sequence whereby the altered polynucleotide retains the biological function or activity of the reference polynucleotide.

20 The term "variant" also includes naturally-occurring allelic variants.

Reference herein to a low stringency includes and encompasses from at least about 0 to at least about 15% v/v formamide and from at least about 1 M to at least about 2 M salt for hybridization, and at least about 1 M to at least about 2 M salt for washing conditions.

25 Generally, low stringency is at from about 25-30°C to about 42°C. The temperature may be altered and higher temperatures used to replace formamide and/or to give alternative stringency conditions. Alternative stringency conditions may be applied where necessary, such as medium stringency, which includes and encompasses from at least about 16% v/v to at least about 30% v/v formamide and from at least about 0.5 M to at least about 0.9 M salt for hybridization, and at least about 0.5 M to at least about 0.9 M salt for washing conditions, or high stringency, which includes and encompasses from at least about 31%

v/v to at least about 50% v/v formamide and from at least about 0.01 M to at least about 0.15 M salt for hybridization, and at least about 0.01 M to at least about 0.15 M salt for washing conditions. In general, washing is carried out T<sub>m</sub> = 69.3 + 0.41 (G+C)% (Marmur and Doty, J. Mol. Biol. 5: 109, 1962). However, the T<sub>m</sub> of a duplex DNA decreases by 1°C with every increase of 1% in the number of mismatch base pairs (Bonner and Laskey, Eur. J. Biochem. 46: 83, 1974). Formamide is optional in these hybridization conditions. Accordingly, particularly preferred levels of stringency are defined as follows: low stringency is 6 x SSC buffer, 0.1% w/v SDS at 25°-42°C; a moderate stringency is 2 x SSC buffer, 0.1% w/v SDS at a temperature in the range 20°C to 65°C; high stringency is 0.1 x SSC buffer, 0.1% w/v SDS at a temperature of at least 65°C.

The present invention extends to recombinant forms of the M-grh molecules as well as derivatives and homologs thereof.

Accordingly, another aspect of the present invention provides an isolated polypeptide having transcription factor activity, said polypeptide comprising a sequence of amino acids encoded by a nucleotide sequence selected from SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or a nucleotide sequence having at least about 60% similarity to any one of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or a nucleotide sequence capable of hybridizing to any one of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or a complementary form thereof under low stringency conditions.

25

In a preferred embodiment, the present invention provides a recombinant M-grh comprising an amino acid sequence selected from SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 or SEQ ID NO:16 or an amino acid sequence having at least about 60% similarity to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 or SEQ ID NO:16.

This aspect of the present invention extends to derivatives, homologs and analogs of M-GRH molecules.

A "derivative" includes a mutant, fragment, part, portion or hybrid molecule. A derivative generally but not exclusively carries a single or multiple amino acid substitution, addition and/or deletion.

A "homolog" includes an analogous polypeptide having at least about 60% similar amino acid sequence from another animal species or from a different locus within the same species.

An "analog" is generally a chemical analog. Chemical analogs of the subject polypeptide contemplated herein include, but are not limited to, modification to side chains, incorporation of unnatural amino acids and/or their derivatives during peptide, polypeptide or protein synthesis and the use of crosslinkers and other methods which impose conformational constraints on the proteinaceous molecule or their analogs.

Examples of side chain modifications contemplated by the present invention include modifications of amino groups such as by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH<sub>4</sub>; amidination with methylacetimidate; acylation with acetic anhydride; carbamoylation of amino groups with cyanate; trinitrobenzylation of amino groups with 2, 4, 6-trinitrobenzene sulphonic acid (TNBS); acylation of amino groups with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation of lysine with pyridoxal-5-phosphate followed by reduction with NaBH<sub>4</sub>.

The guanidine group of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

25

15

The carboxyl group may be modified by carbodiimide activation via O-acylisourea formation followed by subsequent derivitization, for example, to a corresponding amide.

Sulphydryl groups may be modified by methods such as carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulphides with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted maleimide; formation of mercurial derivatives using 4-chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-chloromercuri-4-nitrophenol and other mercurials; carbamoylation with cyanate at alkaline pH.

Tryptophan residues may be modified by, for example, oxidation with N-bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues on the other hand, may be altered by nitration with tetranitromethane to form a 3-nitrotyrosine derivative.

Modification of the imidazole ring of a histidine residue may be accomplished by alkylation with iodoacetic acid derivatives or N-carbethoxylation with diethylpyrocarbonate.

20

25

15

10

Examples of incorporating unnatural amino acids and derivatives during peptide synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl alanine and/or D-isomers of amino acids. A list of unnatural amino acid, contemplated herein is shown in Table 3.

## TABLE 3

	Non-conventional amino acid	Code	Non-conventional amino acid	Code
	α-aminobutyric acid	Abu	L-N-methylalanine	Nmala
	$\alpha$ -amino- $\alpha$ -methylbutyrate	Mgabu	L-N-methylarginine	Nmarg
	aminocyclopropane-	Cpro	L-N-methylasparagine	Nmasn
)	carboxylate		L-N-methylaspartic acid	Nmasp
	aminoisobutyric acid	Aib	L-N-methylcysteine	Nmcys
	aminonorbornyl-	Norb	L-N-methylglutamine	Nmgln
•	carboxylate		L-N-methylglutamic acid	Nmglu
	cyclohexylalanine	Chexa	L-Nmethylhistidine	Nmhis
5	cyclopentylalanine	Cpen	L-N-methylisolleucine	Nmile
	D-alanine	Dal	L-N-methylleucine	Nmleu
	D-arginine	Darg	L-N-methyllysine	Nmlys
	D-aspartic acid	Dasp	L-N-methylmethionine	Nmmet
	D-cysteine	Dcys	L-N-methylnorleucine	Nmnle
0	D-glutamine	Dgln	L-N-methylnorvaline	Nmnva
	D-glutamic acid	Dglu	L-N-methylornithine	Nmorn
	D-histidine	Dhis	L-N-methylphenylalanine	Nmphe
	D-isoleucine	Dile	L-N-methylproline	Nmpro
	D-leucine	Dleu	L-N-methylserine	Nmser
5	D-lysine	Dlys	L-N-methylthreonine	Nmthr
	D-methionine	Dmet	L-N-methyltryptophan	Nmtrp
	D-ornithine	Dom	L-N-methyltyrosine	· Nmtyr
	D-phenylalanine	Dphe	L-N-methylvaline	Nmval
	D-proline	Dpro	L-N-methylethylglycine	Nmetg
30	D-serine	Dser	L-N-methyl-t-butylglycine	Nmtbug
	D-threonine	Dthr	L-norleucine	Nie

	D-tryptophan	Dtrp	L-norvaline	Nva
	D-tyrosine	Dtyr	α-methyl-aminoisobutyrate	Maib
	D-valine	Dval	α-methyl-γ-aminobutyrate	Mgabu
	D-α-methylalanine	Dmala	α-methylcyclohexylalanine	Mchexa
5	D-α-methylarginine	Dmarg	α-methylcylcopentylalanine	Mcpen
	D-α-methylasparagine	Dmasn	α-methyl-α-napthylalanine	Manap
	D-α-methylaspartate	Dmasp	α-methylpenicillamine	Mpen
	D-α-methylcysteine	Dmcys	N-(4-aminobutyl)glycine	Nglu
÷	D-α-methylglutamine	Dmgln	N-(2-aminoethyl)glycine	Naeg
10	D-α-methylhistidine	Dmhis	N-(3-aminopropyl)glycine	Norn
	D-α-methylisoleucine	Dmile	N-amino-α-methylbutyrate	Nmaabu
	D-α-methylleucine	Dmleu	$\alpha$ -napthylalanine	Anap
	D-α-methyllysine	Dmlys	N-benzylglycine	Nphe
	$D$ - $\alpha$ -methylmethionine	Dmmet	N-(2-carbamylethyl)glycine	Ngln
15	D-α-methylornithine	Dmorn	N-(carbamylmethyl)glycine	Nasn
	$D$ - $\alpha$ -methylphenylalanine	Dmphe	N-(2-carboxyethyl)glycine	Nglu
	D-α-methylproline	Dmpro	N-(carboxymethyl)glycine	Nasp
	$D$ - $\alpha$ -methylserine	Dmser	N-cyclobutylglycine	Nebut
	D-α-methylthreonine	Dmthr	N-cycloheptylglycine	Nchep
20	D-α-methyltryptophan	Dmtrp	N-cyclohexylglycine	Nchex
	$D-\alpha$ -methyltyrosine	Dmty	N-cyclodecylglycine	Ncdec
	$D-\alpha$ -methylvaline	Dmval	N-cylcododecylglycine	Ncdod
	D-N-methylalanine	Dnmala	N-cyclooctylglycine	Ncoct
	D-N-methylarginine	Dnmarg	N-cyclopropylglycine	<b>N</b> cpro
25	D-N-methylasparagine	Dnmasn	N-cycloundecylglycine	Nound
	D-N-methylaspartate	Dnmasp	N-(2,2-diphenylethyl)glycine	Nbhm
	D-N-methylcysteine	Dnmcys	N-(3,3-diphenylpropyl)glycine	Nbhe
	D-N-methylglutamine	Dnmgln	N-(3-guanidinopropyl)glycine	Narg
	D-N-methylglutamate	Dnmglu	N-(1-hydroxyethyl)glycine	Nthr
30	D-N-methylhistidine	Dnmhis	N-(hydroxyethyl))glycine	Nser

	D-N-methylisoleucine	Dnmile	N-(imidazolylethyl))glycine	Nhis
	D-N-methylleucine	Dnmleu	N-(3-indolylyethyl)glycine	Nhtrp
	D-N-methyllysine	Dnmlys	N-methyl-γ-aminobutyrate	Nmgabu
	N-methylcyclohexylalanine	Nmchexa	D-N-methylmethionine	Dnmmet
5	D-N-methylornithine	Dnmorn	N-methylcyclopentylalanine	Nmcpen
	N-methylglycine	Nala	D-N-methylphenylalanine	Dnmphe
	N-methylaminoisobutyrate	Nmaib	D-N-methylproline	Dnmpro
	N-(1-methylpropyl)glycine	Nile	D-N-methylserine	Dnmser
	N-(2-methylpropyl)glycine	Nleu	D-N-methylthreonine	Dnmthr
10	D-N-methyltryptophan	Dnmtrp	N-(1-methylethyl)glycine	Nval
	D-N-methyltyrosine	Dnmtyr	N-methyla-napthylalanine	Nmanap
	D-N-methylvaline	Dnmval	N-methylpenicillamine	Nmpen
	γ-aminobutyric acid	Gabu	N-(p-hydroxyphenyl)glycine	Nhtyr
	L-t-butylglycine	Tbug	N-(thiomethyl)glycine	Neys
15	L-ethylglycine	Etg	penicillamine	Pen
	L-homophenylalanine	Hphe	L- $\alpha$ -methylalanine	Mala
	L-α-methylarginine	Marg	$L$ - $\alpha$ -methylasparagine	Masn
	L-α-methylaspartate	Masp	L-α-methyl-t-butylglycine	Mtbug
	L-α-methylcysteine	Mcys	L-methylethylglycine	Metg
20	L-α-methylglutamine	Mgln	L-α-methylglutamate	Mglu
	$L$ - $\alpha$ -methylhistidine	Mhis	$L$ - $\alpha$ -methylhomophenylalanine	Mhphe
	L-α-methylisoleucine	Mile	N-(2-methylthioethyl)glycine	Nmet
	L-α-methylleucine	Mleu	L-α-methyllysine	Mlys
	L-α-methylmethionine	Mmet	L-α-methylnorleucine	Mnle
25	L-α-methylnorvaline	Mnva	$L$ - $\alpha$ -methylomithine	Morn
	L-α-methylphenylalanine	Mphe	L-α-methylproline	Mpro
	L-α-methylserine	Mser	L-α-methylthreonine	Mthr
	L-α-methyltryptophan	Mtrp ·	$L$ - $\alpha$ -methyltyrosine	Mtyr
	L-α-methylvaline	Mval	L-N-methylhomophenylalanine	Nmhphe
3	0 N-(N-(2,2-diphenylethyl)	Nnbhm	N-(N-(3,3-diphenylpropyl)	Nnbhe

10

15

20

25

- 26 -

carbamylmethyl)glycine
1-carboxy-1-(2,2-diphenyl- Nmbc
ethylamino)cyclopropane

carbamylmethyl)glycine

Crosslinkers can be used, for example, to stabilize 3D conformations, using homobifunctional crosslinkers such as the bifunctional imido esters having  $(CH_2)_n$  spacer groups with n=1 to n=6, glutaraldehyde, N-hydroxysuccinimide esters and hetero-bifunctional reagents which usually contain an amino-reactive moiety such as N-hydroxysuccinimide and another group specific-reactive moiety such as maleimido or dithio moiety (SH) or carbodiimide (COOH). In addition, peptides can be conformationally constrained by, for example, incorporation of  $C_{\alpha}$  and  $N_{\alpha}$ -methylamino acids, introduction of double bonds between  $C_{\alpha}$  and  $C_{\beta}$  atoms of amino acids and the formation of cyclic peptides or analogues by introducing covalent bonds such as forming an amide bond between the N and C termini, between two side chains or between a side chain and the N or C terminus.

The present invention further contemplates chemical analogs of the subject polypeptide capable of acting as antagonists or agonists of M-GRH or which can act as functional analogs of M-GRH. Chemical analogs may not necessarily be derived from the instant M-GRH molecules but may share certain conformational similarities. Alternatively, chemical analogs may be specifically designed to mimic certain physiochemical properties of the subject M-GRH molecules. Chemical analogs may be chemically synthesized or may be detected following, for example, natural product screening. The latter refers to molecules identified from various environmental sources such a river beds, coral, plants, microorganisms and insects.

These types of modifications may be important to stabilize the subject M-GRH molecules if administered to an individual or for use as a diagnostic reagent.

30 Other derivatives contemplated by the present invention include a range of glycosylation variants from a completely unglycosylated molecule to a modified glycosylated molecule.

Altered glycosylation patterns may result from expression of recombinant molecules in different host cells.

The present invention also provides a method for identifying a M-GRH, said method comprising screening a nucleotide database and identifying a nucleotide sequence having at least 60% similarity to SEQ ID NO:17 or SEQ ID NO:34 or SEQ ID NO:36 or SEQ ID NO:38 after optimal alignment.

Reference to a "nucleotide database" includes screening an existing genomic or cDNA or mRNA database or screening for a target nucleic acid molecule in a mammalian cell such as using oligonucleotide probes or primers, sequencing the target molecule and comparing the sequence to SEQ ID NO:17 or SEQ ID NO:34 or SEQ ID NO:36 or SEQ ID NO:38.

In an alternative method, a database of mammalian protein sequences is screened for an amino acid sequence having at least 60% similarity to the amino acid sequence encoded by SEQ ID NO:17 or SEQ ID NO:34 or SEQ ID NO:36 or SEQ ID NO:38. Again, a "database" includes a *de novo* protein sequence isolated and identified on a transcription factor isolated form a mammalian cell.

In yet another alternative, a M-grh or its protein product is deemed one which has at least about 60% similarity at the nucleotide level to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or at the amino acid level to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 or SEQ ID NO:15.

Still yet another aspect of the present invention provides a means of identifying a nucleotide sequence likely to encode an M-GRH transcription factor, said method comprising interrogating a mammalian genome database conceptually translated into different reading frames with an amino acid sequence defining *Drosophila GRH* or any one of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 and SEQ ID NO:16 and identifying a nucleotide sequence

corresponding to an amino acid sequence having at least about 60% similarity to *Drosophila GRH* or to any one of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 and SEQ ID NO:16.

Preferably, the genome is conceptually translated into from about 3 to about 6 reading frames and more preferably six reading frames.

It is proposed in accordance with the present invention that the M-GRH transcription factors are involved in the modulation of expression of a number of genes including developmentally regulated genes. Accordingly, aberrations in the M-GRH or M-grh molecules are proposed to cause over or under expression of particular genes leading to a potentially unwanted phenotype. The phenotype may manifest itself pre- or post-natally. A pre-natal manifestation includes at the embryo or fetus stage. Conditions contemplated include developmentally-determined disease conditions such as poor brain development, poor muscle or bone development, aberrations in facial or cranial structures, malformed spinal structures, predispositions to a range of cancers including melanomas and immunological disorders.

Accordingly, another aspect of the present invention contemplates a method for detecting an aberrant phenotype or a propensity for an aberrant phenotype to develop, said method comprising screening for a variation in a nucleotide sequence encoding a mammalian MGR, BOM and/or SOM or their homologs.

Reference herein to "MGR", "BOM" and "SOM" includes murine and human forms of these molecules such as human MGR p49 (SEQ ID NO:2), human MGR p70 (SEQ ID NO:4), human BOM (SEQ ID NO:6), human SOM (SEQ ID NO:8), murine MGR p61 (SEQ ID NO:10), murine MGR p70 (SEQ ID NO:12), murine BOM (SEQ ID NO:14) and murine SOM (SEQ ID NO:16).

A homolog of MGR, BOM and SOM is as herein defined including a molecule having at least about 60% amino acid sequence similarity to MGR, BOM or SOM or at least about

60% nucleic acid similarity to mgr, bom or som or a nucleic acid molecule capable of hybridizing to the coding strands of mgr, bom or som or complementary forms thereof under low stringency conditions.

Aberrations may also be detectable at the amino acid level when the mammalian homologs of *Drosophila grh* encode protein transcription factors.

Accordingly, another aspect of the present invention contemplates a method for detecting an aberrant phenotype or a propensity for an aberrant phenotype to develop, said method comprising screening for a variation in an amino acid sequence encoding MGR, BOM and/or SOM or their homologs.

As above, reference to MGR, BOM and SOM include amino acid sequences defining human MGR p49 (SEQ ID NO:2), human MGR p70 (SEQ ID NO:4), human BOM SEQ ID NO:6), human SOM (SEQ ID NO:8), murine MGR p61 (SEQ ID NO:10), murine MGR p70 (SEQ ID NO:12), murine BOM (SEQ ID NO:14) and murine SOM (SEQ ID NO:16).

As stated above, the mammalian transcription factors and their genetic sequences have a range of diagnostic and therapeutic utilities. The detection of an aberrant transcription factor or a nucleotide sequence encoding an aberrant transcription factor is indicative of a disease condition including a degenerative or developmental disease condition.

Any number of methods may be employed to detect aberrant transcription factors or their genetic sequences. Immunological testing is one particular method. Accordingly, the present invention extends to antibodies and other immunological agents directed to or preferably specific for the mammalian transcription factors or a fragment thereof. The antibodies may be monoclonal or polyclonal or may comprise Fab fragments or synthetic forms.

20

25

30

Specific antibodies can be used to screen for the subject mammalian transcription factors and/or their fragments. Techniques for the assays contemplated herein are known in the art and include, for example, sandwich assays and ELISA.

It is within the scope of this invention to include any second antibodies (monoclonal, polyclonal or fragments of antibodies or synthetic antibodies) directed to the first mentioned antibodies referred to above. Both the first and second antibodies may be used in detection assays or a first antibody may be used with a commercially available anti-immunoglobulin antibody. An antibody as contemplated herein includes any antibody specific to any region of the mammalian transcription factors.

Both polyclonal and monoclonal antibodies are obtainable by immunization with mammalian transcription factors or antigenic fragments thereof and either type is utilizable for immunoassays. The methods of obtaining both types of sera are well known in the art. Polyclonal sera are less preferred but are relatively easily prepared by injection of a suitable laboratory animal with an effective amount of subject polypeptide, or antigenic parts thereof, collecting serum from the animal and isolating specific sera by any of the known immunoadsorbent techniques. Although antibodies produced by this method are utilizable in virtually any type of immunoassay, they are generally less favoured because of the potential heterogeneity of the product.

The use of monoclonal antibodies in an immunoassay is particularly preferred because of the ability to produce them in large quantities and the homogeneity of the product. The preparation of hybridoma cell lines for monoclonal antibody production derived by fusing an immortal cell line and lymphocytes sensitized against the immunogenic preparation can be done by techniques which are well known to those who are skilled in the art.

Another aspect of the present invention contemplates, therefore, a method for detecting a mammalian transcription factor or fragment thereof in a biological sample from a subject, said method comprising contacting said biological sample with an antibody specific for said mammalian transcription factor or fragment thereof or its derivatives or homologs for

a time and under conditions sufficient for an antibody-polypeptide complex to form, and then detecting said complex.

A biological sample includes a cell extract.

5

15

25

Reference to a "mammalian transcription factor" is considered to be a reference to a homolog of Drosophia grh, i.e. M-GRH.

The presence of the instant mammalian transcription factors or their fragments may be detected in a number of ways such as by Western blotting and ELISA procedures. A wide range of immunoassay techniques are available as can be seen by reference to U.S. Patent Nos. 4,016,043, 4,424,279 and 4,018,653.

Sandwich assays are among the most useful and commonly used assays and are favoured for use in the present invention. A number of variations of the sandwich assay technique exist, and all are intended to be encompassed by the present invention. Briefly, in a typical forward assay, an unlabeled antibody is immobilized on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen 20 complex, a second antibody specific to the antigen, labeled with a reporter molecule capable of producing a detectable signal is then added and incubated, allowing time sufficient for the formation of another complex of antibody-antigen-labeled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal produced by the reporter molecule. The results may either be qualitative, by simple observation of the visible signal, or may be quantitated by comparing with a control sample containing known amounts of hapten. Variations on the forward assay include a simultaneous assay, in which both sample and labeled antibody are added simultaneously to the bound antibody. These techniques are well known to those skilled in the art, including any minor variations as will be readily apparent. In accordance with the present invention the sample is one which might contain a subject transcription factor including by tissue biopsy, blood, synovial fluid and/or lymph. The sample is,

therefore, generally a biological sample comprising biological fluid. The transcription factor is likely to be in blood or other fluid in the case where cell apoptosis is occurring.

In the typical forward sandwich assay, a first antibody having specificity for the instant polypeptide or antigenic parts thereof, is either covalently or passively bound to a solid surface. The solid surface is typically glass or a polymer, the most commonly used polymers being cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene. The solid supports may be in the form of tubes, beads, discs of microplates, or any other surface suitable for conducting an immunoassay. The binding processes are well-known in the art and generally consist of cross-linking covalently binding or 10 physically adsorbing, the polymer-antibody complex is washed in preparation for the test sample. An aliquot of the sample to be tested is then added to the solid phase complex and incubated for a period of time sufficient (e.g. 2-40 minutes or where more convenient, overnight) and under suitable conditions (e.g. for about 20°C to about 40°C) to allow binding of any subunit present in the antibody. Following the incubation period, the 15 antibody subunit solid phase is washed and dried and incubated with a second antibody specific for a portion of the hapten. The second antibody is linked to a reporter molecule which is used to indicate the binding of the second antibody to the hapten.

An alternative method involves immobilizing the target molecules in the biological sample and then exposing the immobilized target to specific antibody which may or may not be labeled with a reporter molecule. Depending on the amount of target and the strength of the reporter molecule signal, a bound target may be detectable by direct labelling with the antibody. Alternatively, a second labeled antibody, specific to the first antibody is exposed to the target-first antibody complex to form a target-first antibody-second antibody tertiary complex. The complex is detected by the signal emitted by the reporter molecule.

By "reporter molecule" as used in the present specification, is meant a molecule which, by its chemical nature, provides an analytically identifiable signal which allows the detection of antigen-bound antibody. Detection may be either qualitative or quantitative. The most commonly used reporter molecules in this type of assay are either enzymes, fluorophores

15

or radionuclide containing molecules (i.e. radioisotopes) and chemiluminescent molecules. In the case of an enzyme immunoassay, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. As will be readily recognized, however, a wide variety of different conjugation techniques exist, which are readily available to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase and alkaline phosphatase, amongst others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable colour change. Examples of suitable enzymes include alkaline phosphatase and peroxidase. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. In all cases, the enzyme-labeled antibody is added to the first antibody hapten complex, allowed to bind, and then the excess reagent is washed away. A solution containing the appropriate substrate is then added to the complex of antibody-antigenantibody. The substrate will react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of hapten which was present in the sample. "Reporter molecule" also extends to use of cell agglutination or inhibition of agglutination such as red blood cells on latex beads, and the like.

Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be 20 chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labeled antibody adsorbs the light energy, inducing a state to excitability in the molecule, followed by emission of the light at a characteristic colour visually detectable with a light microscope. 25 The fluorescent labeled antibody is allowed to bind to the first antibody-hapten complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to the light of the appropriate wavelength the fluorescence observed indicates the presence of the hapten of interest. Immunofluorescene and EIA techniques are both very well established in the art and are particularly preferred for the present method. However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules, 30 may also be employed.

The present invention also contémplates genetic assays such as involving PCR analysis to detect RNA expression products of a genetic sequence encoding a mammalian transcription factor. The genetic assays may also be able to detect nucleotide polymorphisms or other substitutions, additions and/or deletions in the nucleotide sequence of a mammalian transcription factor. Changes in levels of mammalian transcription factor expression such as following mutations in the promoter or regulatory regions or loss of mammalian transcription factor activity following mutations in mammalian transcription factor nucleotides is proposed to be indicative of a disease condition or a propensity for a disease condition to develop. For example, a cell biopsy could be obtained and DNA or RNA extracted. Alternative methods which may be used alone or in conjunction with other methods include direct nucleotide sequencing or mutation scanning such as single stranded conformation polymorphoms analysis (SSCP) as well as specific oligonucleotide hybridization, denaturing high performance liquid chromatography, first nucleotide change (FNC) amongst others. 15

The present invention extends to polymorphisms which in the M-grh genes leads to healthy or abnormal phenotypes.

The present invention further contemplates kits to facilitate the rapid detection of 20 mammalian transcription factors or their fragments in a subject's biological fluid.

Again, a biological fluid includes a cell extract such as a DNA/RNA extract.

- Still yet another aspect of the present invention contemplates genomic sequences including 25 gene sequences encoding a mammalian transcription factor as well as regulatory regions such as promoters, terminators and transcription/translation enhancer regions associated with the gene encoding a mammalian transcription factor.
- The term "gene" is used in its broadest sense and includes cDNA corresponding to the 30 exons of a gene. Accordingly, reference herein to a "gene" is to be taken to include:-

5

- (i) a classical genomic gene consisting of transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (i.e. introns, 5'- and 3'- untranslated sequences); or
- (ii) mRNA or cDNA corresponding to the coding regions (i.e. exons) and 5'- and 3'- untranslated sequences of the gene.

The term "gene" is also used to describe synthetic or fusion molecules encoding all or part of an expression product. In particular embodiments, the term "nucleic acid molecule" and "gene" may be used interchangeably.

In a particularly useful embodiment, the present invention provides a promoter for the mammalian transcription factor gene. The identification of the promoter permits developmentally-regulated expression of particular genetic sequences. The latter would include a range of therapeutic molecules such as cytokines, growth factors, antibiotics or other molecules to assist in the treatment of particular disease conditions.

Accordingly, another aspect of the present invention provides a M-grh-specific promoter or functional derivative or homolog thereof, said promoter in situ operably linked to a nucleotide sequence comprising any one of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or their complementary forms or a nucleotide sequence having at least about 60% similarity to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or their complementary forms or a nucleotide sequence capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or their complementary forms under low stringency conditions.

30 The promoter is conveniently resident in a vector which comprises unique restriction sites to facilitate the introduction of genetic sequences operably linked to the promoter.

20

25

All such constructs are useful in order to produce recombinant M-GRH molecules and/or in gene therapy protocols.

5 The present invention further contemplates a genetically modified animal.

More particularly, the present invention provides an animal model useful for screening for agents capable of ameliorating the effects of an aberrant M-GRH or M-grh gene. In one embodiment, the animal model produces low amounts of M-grh. Such an animal would have a predisposition for a range of diseases including developmentally regulated diseases. The animal model is useful for screening for agents which ameliorate such conditions.

Accordingly, another aspect of the present invention provides a genetically modified animal wherein said animal produces low amounts of M-grh relative to a non-genetically modified animal of the same species. Reference to "low amounts" includes zero amounts or up to about 10% lower than normalized amounts.

Preferably, the genetically modified animal is a mouse, rat, guinea pig, rabbit, pig, sheep or goat. More preferably, the genetically modified animal is a mouse or rat. Most preferably, the genetically modified animal is a mouse.

Accordingly, a preferred aspect of the present invention provides a genetically modified mouse wherein said mouse produces low amounts of M-grh relative to a non-genetically modified mouse of the same strain.

The animal model contemplated by the present invention comprises, therefore, an animal which is substantially incapable of producing a M-grh. Generally, but not exclusively, such an animal is referred to as a homozygous or heterozygous M-grh-knockout animal.

The animal models of the present invention may be in the form of the animals or may be, for example, in the form of embryos for transplantation. The embryos are preferably maintained in a frozen state and may optionally be sold with instructions for use.

The genetically modified animals may also produce larger amounts of M-GRH For example, over expression of normal M-grh or mutant M-grh may produce dominant negative effects and may become useful disease models.

Accordingly, another aspect of the present invention is directed to a genetically modified animal over-expressing genetic sequences encoding M-grh.

A genetically modified animal includes a transgenic animal, or a "knock-out" or "knock-in" animal.

15 Yet another aspect of the present invention provides a targeting vector useful for inactivating a gene encoding M-GRH, said targeting vector comprising two segments of genetic material encoding said M-GRH flanking a positive selectable marker wherein when said targeting vector is transfected into embryonic stem (ES) cells and the marker selected, an ES cell is generated in which the gene encoding said M-GDH is inactivated by homologous recombination.

Preferably, the ES cells are from mice, rats, guinea pigs, pigs, sheep or goats. Most preferably, the ES cells are from mice.

25 Still yet another aspect of the present invention is directed to the use of a targeting vector as defined above in the manufacture of a genetically modified animal substantially incapable of producing M-GRH.

Even still another aspect of the present invention is directed to the use of a targeting vector as defined above in the manufacture of a genetically modified mouse substantially incapable of producing M-GRH.

Preferably, the vector is DNA. A selectable marker in the targeting vector allows for selection of targeted cells that have stably incorporated the targeting DNA. This is especially useful when employing relatively low efficiency transformation techniques such as electroporation, calcium phosphate precipitation and liposome fusion where typically fewer than 1 in 1000 cells will have stably incorporated the exogenous DNA. Using high efficiency methods, such as microinjection into nuclei, typically from 5-25% of the cells will have incorporated the targeting DNA; and it is, therefore, feasible to screen the targeted cells directly without the necessity of first selecting for stable integration of a selectable marker. Either isogenic or non-isogenic DNA may be employed.

Examples of selectable markers include genes conferring resistance to compounds such as antibiotics, genes conferring the ability to grow on selected substrates, genes encoding proteins that produce detectable signals such as luminescence. A wide variety of such markers are known and available, including, for example, antibiotic resistance genes such as the neomycin resistance gene (neo) and the hygromycin resistance gene (hyg). Selectable markers also include genes conferring the ability to grow on certain media substrates such as the tk gene (thymidine kinase) or the hprt gene (hypoxanthine phosphoribosyltransferase) which confer the ability to grow on HAT medium (hypoxanthine, aminopterin and thymidine); and the bacterial gpt gene (guanine/xanthine phosphoribosyltransferase) which allows growth on MAX medium (mycophenolic acid, adenine and xanthine). Other selectable markers for use in mammalian cells and plasmids carrying a variety of selectable markers are described in Sambrook et al., Molecular Cloning - A Laboratory Manual, Cold Spring Harbour, New York, USA, 1990.

25

30

20

10

The preferred location of the marker gene in the targeting construct will depend on the aim of the gene targeting. For example, if the aim is to disrupt target gene expression, then the selectable marker can be cloned into targeting DNA corresponding to coding sequence in the target DNA. Alternatively, if the aim is to express an altered product from the target gene, such as a protein with an amino acid substitution, then the coding sequence can be

25

30

modified to code for the substitution, and the selectable marker can be placed outside of the coding region, for example, in a nearby intron.

The selectable marker may depend on its own promoter for expression and the marker gene may be derived from a very different organism than the organism being targeted (e.g. prokaryotic marker genes used in targeting mammalian cells). However, it is preferable to replace the original promoter with transcriptional machinery known to function in the recipient cells. A large number of transcriptional initiation regions are available for such purposes including, for example, metallothionein promoters, thymidine kinase promoters, β-actin promoters, immunoglobulin promoters, SV40 promoters and human cytomegalovirus promoters. A widely used example is the pSV2-neo plasmid which has the bacterial neomycin phosphotransferase gene under control of the SV40 early promoter and confers in mammalian cells resistance to G418 (an antibiotic related to neomycin). A number of other variations may be employed to enhance expression of the selectable markers in animal cells, such as the addition of a poly(A) sequence and the addition of synthetic translation initiation sequences. Both constitutive and inducible promoters may be used.

The DNA is preferably modified by homologous recombination. The target DNA can be in any organelle of the animal cell including the nucleus and mitochondria and can be an intact gene, an exon or intron, a regulatory sequence or any region between genes.

Homologous DNA is a DNA sequence that is at least 70% identical with a reference DNA sequence. An indication that two sequences are homologous is that they will hybridize with each other under stringent conditions (Sambrook et al., 1990, supra).

The present invention further contemplates co-suppression (i.e. sense suppression) and antisense suppression to down-regulate expression of M-grh This would generally occur in a target test animal such as to generate a disease model.

In addition to providing a diagnostic capability as described above, the isolated nucleic

15

25

30

acid molecules of the present invention may also provide a therapeutic capability by being used to correct or complement an abnormality detected in a subject. To deliver the appropriate sequence to a recipient cell or tissue of a subject, an isolated nucleic acid molecule of the present invention may be cloned into a suitable genetic construct such as a suitable vector.

Accordingly, a further aspect of the present invention contemplates a genetic construct comprising a nucleotide sequence encoding an M-grh selected from SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or a variant thereof or a nucleotide sequence having at least 60% similarity to one or more of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 or SEQ ID NO:15 or a variant thereof or a nucleotide sequence capable of hybridizing to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:15 under low stringency conditions or a variant thereof or a complementary form thereof.

A "vector" is a polynucleotide molecule, preferably a DNA molecule derived, for example, from a plasmid, bacteriophage, or plant virus, into which a polynucleotide can be inserted or cloned. A vector preferably contains one or more unique restriction sites and can be capable of autonomous replication in a defined host cell including a target cell or tissue or a progenitor cell or tissue thereof, or be integrable with the genome of the defined host such that the cloned sequence is reproducible. Accordingly, the vector may be an autonomously replicating vector, i.e. a vector that exists as an extra-chromosomal entity, the replication of which is independent of chromosomal replication. Examples include a linear or closed circular plasmid, an extra-chromosomal element, a mini-chromosome, or an artificial chromosome. The vector may also contain a means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. A vector system may comprise a single vector or plasmid, two or more vectors or plasmids, which together contain the total DNA to be introduced into the genome of the host cell, or a transposon.

Vectors suitable for gene therapy applications are well known in the art. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which it is to be introduced. The vector may also include an additional genetic construct comprising a selection marker such as an antibiotic resistance gene that can be used for selection of suitable transformants. Examples of such resistance genes are known to those skilled in the art and include the *nptII* gene that confers resistance to the antibiotics kanamycin, and G418 (Geneticin®) and the *hph* gene which confer resistance to the antibiotic hygromycin B.

10

Accordingly, in a related embodiment, the present invention provides a genetic construct comprising a promoter or functional equivalent thereof operably linked to a nucleotide sequence of the invention.

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences of a classical genomic gene, which is required for accurate transcription initiation, with or without a CCAAT box sequence and additional regulatory elements (i.e. upstream activating sequences, enhancers and silencers), which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. A promoter is usually, but not necessarily, positioned upstream (5') of a gene region, the expression of which it regulates. Furthermore, the regulatory elements comprising a promoter are usually positioned within 2 kb of the start site of transcription of the gene. As is known in the art, some variation in this distance can be accommodated without loss of promoter function.

25

The selection of an appropriate promoter sequence to regulate expression of a transcription factor encoded by an isolated nucleic acid molecule of the present invention is an important consideration. Examples of suitable promoters include viral, fungal, bacterial, animal and plant derived promoters capable of functioning in eukaryotic animal cells and, especially, human cells. The promoter may regulate the expression of the nucleic acid molecule differentially with respect to the cell, tissue or organ in which expression occurs,

or with respect to the developmental stage at which expression occurs.

Preferably, the promoter is capable of regulating expression of a nucleic acid molecule in a eukaryotic cell, tissue or organ, at least during the period of time over which the regulated gene is expressed therein, and more preferably also immediately preceding the commencement of detectable expression of the regulated gene in said cell, tissue or organ.

Particularly preferred promoters for use with the nucleic acid molecules of the present invention include the bacteriophage T7 promoter, bacteriophage T3 promoter, SP6 promoter, *lac* operator-promoter, *tac* promoter, SV40 late promoter, SV40 early promoter, RSV-LTR promoter, CMV IE promoter, CaMV 35S promoter, SCSV promoter, SCBV promoter and the like. Those skilled in the art will readily be aware of additional promoter sequences other than those specifically described.

- In the present context, the terms "in operable connection with" or "operably linked" or similar shall be taken to indicate that expression of the nucleic acid molecule is under the control of the promoter sequence, with which it is spatially connected, in a cell, tissue, organ or whole organism.
- The genetic construct of the present invention may also comprise a 3' non-translated sequence. A 3' non-translated sequence refers to that portion of a gene comprising a DNA segment that contains a polyadenylation signal and any other regulatory signals capable of effecting mRNA processing or gene expression. The polyadenylation signal is characterized by effecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognized by the presence of homology to the canonical form 5' AATAAA-3' although variations are not uncommon.

Accordingly, a genetic construct comprising a nucleic acid molecule of the present invention, operably linked to a promoter, may be cloned into a suitable vector for delivery to a cell or tissue in which regulation is faulty, malfunctioning or non-existent, in order to rectify and/or provide the appropriate regulation. Vectors comprising appropriate genetic

constructs may be delivered into target eukaryotic cells by a number of different means well known to those skilled in the art of molecular biology.

The present invention further contemplates the use of an M-GRH or M-grh in the manufacture of a medicament for the treatment of a disease condition in a mammal such as a human.

The present invention is further directed to promoters and 3'- and 5'-regulatory regions associated with genomic forms of M-grh genes. These regions can be readily identified by, for example, chromosome walking using M-grh nucleic acid molecules or probes or primers therefrom.

The present invention is further described by the following non-limiting Examples.

#### **EXAMPLE 1**

#### Polymerase chain reaction

For RT-PCR, first strand cDNA was prepared from 2 µg of mRNA from primary tissues using random hexamers. Each cDNA sample was appropriately diluted to give similar amplification of S14 RNA under the same PCR conditions. The primer sequences are detailed below. The PCR conditions were 94°C for 2 min followed by 35 cycles of 94°C for 30 sec, 60°C for 30 sec and 72°C for 45 sec with a final extension at 72°C for 5 min. All PCR products were electrophoresed on 1.5% w/v agarose gels, transferred to nitrocellulose and analyzed by Southern blot using <sup>32</sup>P-radiolabeled internal oligonucleotides as probes. Membranes were then autoradiographed for 2 hr at -70°C.

The following primers were used to amplify probes for cDNA library screening and for RT-PCR:-

[SEQ ID NO:25]

[SEQ ID NO:26]

15

#### human p49 mgr

	numan p49 mgr	
	5'-GAAGTCTTTGATGCCCTGATG-3'	[SEQ ID NO:19]
	5'-AACCCATTCCCTCGACATAGA-3'	[SEQ ID NO:20]
20	human p70 mgr	
	5-AGCGCGATGACACAGGAGTA-3'	[SEQ ID NO:21]
	5'-CGTTGCTATGGAGACAGTGA-3'	[SEQ ID NO:22]
	human bom	
25	5'-CCGTTTAACAAGGACACTGC-3'	[SEQ ID NO:23]
	5'-CTGGAAGCCACCAAATCTCT-3'	[SEQ ID NO:24]
	murine p70 mgr	

5'-AGCGCGATGACACAGGAGTA-3'

5'-AGTGCCAGAGCTGAACTGAT-3'

#### murine p61 mgr

5'-TCCATGGGTTCCTTGAGTTC-3' [SEQ ID NO:27] 5'-AGTGCCAGAGCTGAACTGAT'-3' [SEQ ID NO:28]

#### 5 murine bom

5'-AAAGGGGAGCGAGTTCATTG-3' [SEQ ID NO:29] 5'-AGAGCTCTCGGTGATGGATA-3' [SEQ ID NO:30]

10

#### **EXAMPLE 2**

#### Cloning of human and murine mgr and bom

Human p49 mgr was cloned from a fetal brain cDNA phage library in the λZAP II vector (Stratagene). The cDNA encoding the longer human MGR isoform was amplified by RTPCR from human kidney mRNA. The cDNA encoding the smaller murine isoform of MGR was cloned from a 17.5-day embryo phage library in the Lambda TripelEx vector (Clontech). The murine p70 cDNA was amplified from murine kidney mRNA by RT-PCR. The human bom cDNA was isolated form a placental phage library in the Lambda ZAP II vector (Stratagene) and the murine cDNA from an embryonic carcinoma cell line (P19) phage library in the Uni-ZAP XR vector (Stratagene). The murine MGR genomic locus was obtained from a 129SVJ phage library in the Lambda FIX II vector (Stratagene).

From similarity searches of GenBank databases, using the GRH protein sequence as a query, two murine expressed sequence tag (EST) entries were found from adult brain and ovary and one human EST entry from fetal brain that were not identical to any previously reported genes, yet shared high degrees of homology with each other and grh. These sequences were used to design murine and human primers and amplified probes from murine adult brain and ovary and human adult brain cDNA. The murine probe from adult brain cDNA was used in a screen of a day 17.5 mouse embryo cDNA library to obtain a full length clone of a gene referred to as mammalian grainyhead (mgr) due to its sequence and functional homology and similar expression pattern to that of the fly gene. The human

probe derived from adult brain cDNA was used to obtain a full length cDNA clone from a human fetal brain library. Amino acid sequence comparison reveals this to be the human homolog of MGR with 94% identity at the amino acid level.

- The murine probe derived from ovary cDNA was used in a screen of a murine teratocarcinoma cell line (P19) cDNA library to obtain a full length clone of a novel gene distinct from but highly related to mgr named brother-of-mgr (bom). The homology between mgr and bom suggests that mgr and bom arose through gene duplication.
- 10 The human homolog of bom was obtained using primers derived from a high throughput genome sequencing (HTGS) database entry with homology to murine bom. These were used to amplify a probe from a human placental cDNA library that was then screened to yield a full length human cDNA clone. Amino acid sequence comparison between murine and human BOM revealed 94% identity.

The sequence alignments between grh, mgr, bom, CP2 and LBP-1a revealed that mgr and bom are more closely related to grh than the previously identified homologs CP2 and LBP-. 1a (Table 4). This homology is particularly evident in the DNA binding and dimerization domains emphasizing the importance of protein/protein and protein/DNA interactions for the function of these factors.

20

15

TABLE 4 Amino acid sequence comparison of GRH-like genes and Drosophila grh

Amino acid identity/ similarity to Grainyhead (%)	Overall	DNA-binding domain	Dimerization domain
MGR	37/52	48/64	39/61
BOM	35/52	46/63	37/61
SOM	33/48	42/60	38/57
CP2	26/42	32/52	29/47
LBP-1a	23/39	31/51	28/43

# EXAMPLE 3 Identification of a second isoform of MGR

A striking feature of the alignment between MGR and BOM was the absence of an MGR domain corresponding to the first 93 amino acids of BOM. In view of the absence of tissue-specific isoforms of GRH, the EST database was searched for similar sequences using the 5' end of bom as a query. A highly similar but non-identical sequence in an EST from murine kidney was located. The most 3' 30 nucleotides of this EST was identical to 30 nucleotides close to the 5' end of the mgr. Based on this, primers were designed from the kidney EST and mgr cDNA sequences and amplified a product of the predicted size from murine kidney cDNA. A similar product was also amplified from human kidney cDNA. Amino acid sequence analysis of the murine product revealed that it was highly homologous to the 5' end of the BOM protein and contiguous with the mgr open reading frame. However, it lacked the first 11 amino acids of a previously isolated mgr clone suggesting the presence of alternate splicing. To examine this, the murine mgr genomic locus was isolated and mapped. As shown in Figure 1B, the first three coding exons in the locus are exclusive to the p70 isoform of mgr. In contrast, the shorter isoform of mgr's (p61) first coding exon is absent in the p70 isoform. Significantly, the 5' end of this exon lacks a splice acceptor site explaining its absence from the longer isoform. Instead, promoter sequences with a clear TATA box and CAP site are evident in close proximity to the translation initiation site (Figure 1C). Subsequent mapping of the human genomic locus revealed that murine exon four was conserved in the human p70 protein but was absent in

5

15

20

25

- 48 -

the 49 kDa isoform of MGR.

#### **EXAMPLE 4**

## The first three exons of the mgr genomic locus encode transcriptional activation domain

Although significant sequence homology exists between grh and the shorter mgr isoforms and p70 mgr, the isoleucine rich transcriptional activation domain identified in the fly protein is not conserved. Examination of the MGR-coding sequences failed to reveal a region homologous to other known transactivation domains. In view of the high degree of conservation of the first three coding exons of p70 mgr and bom, it was postulated that this could be the functional domain responsible for activation. To address this, the cDNA fragment encoding the first 93 amino acids of human p70 MGR (encoded by the first three exons) was subcloned in frame with the GAL4 DNA binding domain in a mammalian expression vector. The comparable region of BOM and the full length p49 MGR cDNA in frame into this vector was also cloned. These plasmids were co-transfected into the human 293T cell line with a reporter plasmid containing five concatamerized GAL4 DNA binding sites upstream of the chloramphenicol acetyltransferase (CAT) gene. The vector containing only the GALA DNA-BD or containing the VP16 activation domain fused to the GAL DNA-BD served as the negative and positive controls, respectively. As shown in Figure 3, transcriptional activation of the CAT gene was observed with VP16, p70 MGR and the bom containing plasmids. No activation was observed with p49 mgr or the empty vector. These findings confirm the presence of a highly conserved activation domain in the p70 mgr and bom that is lacking in p49 mgr.

25

15


20

5

#### **EXAMPLE 5**

#### MGR binds to known GRH binding sites

To determine the extent of the functional homology between GRH and MGR, it was initially examined whether the mammalian protein could bind to the well-characterized binding sites for the *Drosophila* factor in the *Dopa decarboxylase* and *PCNA* gene



20

25

10

As shown in Figure 2A, a specific protein/DNA complex was observed with the *PCNA* probe in the presence of pre-immune sera (lanes 1 and 3). This complex was supershifted with the addition of anti-p70 specific antisera raised against peptides in the amino terminal region of the protein (lane 4) and ablated with the addition of anti-MGR antisera raised against peptides common to p49 and p70 MGR in the dimerization domain of the protein (lane 2). Neither antisera cross-reacted with BOM. Similar results were obtained with the *Dopa decarboxylase* promoter probe (Figure 2B).

#### **EXAMPLE 6**

#### MGR binds to the human Engrailed-1 promoter

Many Drosophila genes regulated by GRH have known mammalian homologs. In terms of functional homology, Engrailed-1 (En-1) is one of the bests characterized. The En-1 promoter was, therefore, examined for the grainyhead consensus DNA binding sequence derived from a comparison of the Drosophila Ultrabithorax, Dopa decarboxylase and fushi tarazu promoters (Dynlacht et al., Genes Dev. 3: 1677-1688, 1989). As shown in Figure 3A, a highly conserved region was identified in the proximal En-1 promoter. Moreover, this sequence was also largely conserved in the DNAseI footprint attributed to grh in the Drosophila engrailed promoter (Soeller et al., Genes Dev. 2: 68-81, 1988). The ability of this region of the human En-1 promoter to compete off MGR binding to the Ddc probe (Figure 3B) in an EMSA with nuclear extract from JEG-3 cells was examined. As shown in Figure 3B, the specific MGR/DNA complex observed with the Ddc probe (lane 15 1) was supershifted with the addition of MGR antisera 67 (lane 2) and ablated with the addition of a 50-fold excess of unlabeled Ddc probe as competitor (lane 3). Addition of a 10- (lane 4) or 20-fold (lane 5) excess of unlabeled En-1 probe also markedly reduced the binding of MGR to the Ddc probe.

20

25

#### **EXAMPLE 7**

#### MGR activates transcription

To determine the functional significance of this binding, this region of the En-1 promoter was linked to a minimal globin gene promoter/luciferase reporter gene construct and transfected it into the MGR null cell line COS, in the presence of p70 MGR mammalian expression vector or the empty vector. Transfection of the minimal promoter/reporter or the TK promoter linked to a Renilla luciferase gene with either vector served as the controls. As shown in Figure 3C, expression of p70 MGR dramatically enhanced the transcriptional activity of the En-1 promoter (solid bars) but not the control minimal promoter (open bars) or the TK promoter (hatched bars).

#### **EXAMPLE 8**

#### Cloning of full-length human SOM

Human SOM was cloned using primers derived from a high through-put genomic sequence (HTGS) and a human expression sequence tag (EST) obtained from GenBank databases which, respectively, aligned with the dimerization domain and the activation domain of other MGR members. Using nested RT-PCR and human tonsil cDNA, another contig spanning 1300 nucleotides was obtained.

10 Utilizing 5' RACE, further oligoprimers and human testis cDNA, a 210 nucleotide sequence incorporating the initiating ATG was obtained. A contig of these overlapping sequences revealed the full length human SOM which upon alignment with other existing MGR family members showed >60% similarity at the protein level with conservation at the 5' activation, DNA-binding and dimerization domains.

15

#### **EXAMPLE 9**

#### Cloning of full-length murine SOM

A murine EST (GenBank) from optic cup tissue was identified, which when aligned with other murine homologs of the MGR family showed 70% similarity at the amino acid level, in the region of the DNA binding domain. Using semi-nested RT-PCR with murine testis cDNA, a 286 nucleotide sequence was amplified, cloned and sequenced for use as a probe.

Subsequently, a murine brain cDNA library (Stratagene) was screened. One clone was taken through to quaternary stage. This clone was excised from λZAP II vector into pBluescript and sequenced in both directions. A 1200 nucleotide length sequence was obtained, whichi lacked the 5' end. This was subsequently identified using 5' RACE from murine testis cDNA. A contig of these two sequences revealed the full length murine SOM.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

#### **BIBLIOGRAPHY**

Altschul et al., Nucl. Acids Res. 25: 3389, 1997.

Attardi and Tjian, Genes Dev. 7: 1341-1353, 1993;

Ausubel et al. In Current Protocols in Molecular Biology, John Wiley & Sons Inc, 1994-1998, Chapter 15.

Biggin and Tjian, Cell 53: 699-711, 1988;

Bray and Kafatos, Genes Dev. 5: 1672-1683, 1991;

Bray et al., Genes Dev. 3: 1130-1145, 1989;

Bucher, J. Mol. Biol. 212: 563-578, 1990;

Dynlacht et al., Cell 56: 563-576, 1991;

Dynlacht et al., Genes Dev. 3: 1677-1688, 1989;

Dynlacht et al., Genes Dev. 3: 1677-1688, 1989;

Hayashi et al., J. Biol. Chem. 274: 35080-35088, 1999;

Huang et al., Genes Dev. 9: 3177-3189, 1995;

Jane et al., EMBO J. 14: 97-105, 1995;

Liaw et al., Genes Dev. 9: 3163-3176, 1995;

Logan et al., Dev. Genet. 13: 345-358, 1992;

Ramamurthy et al., J. Biol. Chem. 276: 7836-7842, 2001;

Soeller et al., Genes Dev. 2: 68-81, 1988;

Soeller et al., Genes Dev. 2: 68-81, 1988;

Sueyoshi et al., Mol. Cell Biol. 15: 4158-4166, 1995;

Uv et al., Mol. Cell Biol. 14: 4020-4031, 1994;

Uv et al., Mol. Cell Biol. 17: 6727-6735, 1997;

Uv et al., Mol. Cell. Biol. 17: 6727-6735, 1997;

Volker et al., Genes Development 11: 1435-1446, 1997;


Zhou et al., Mol. Cell Biol. 20: 7662-7672, 2000.

```
PTO MGR: 1: DYPHICISOR

PTO MGR: 1: DYPHICISOR

1 NOTE THAT THE PROPERTY OF TH
```

### Figure 1A



## Figure 1B

ABGGGAGTAZ<u>ICAGTCTTT</u>CCPCCATGGCTTTCTTTAGTTTCTCLATATGGCCTTCCTTGATGATGATGATGATGTTGT CAP 05to T

Figure 1C




Figure 2A




Figure 2B

TAGGAGTGAGCCGGCGAAACCGGTTCGCATGGGGCAGGTGACA Drosophila engrailed
ACACCCACCCACAAACACAAACCGGCAGTGACAACAACCACCCAT human engrailed-1
CAAACCGGT TG
YNAACYGGTYYTGCGG Drosophila consensus

## Figure 3A




Figure 3B




Figure 3C

- 1 -

### SEQUENCE LISTING

<110>	Melbo	ourne	e Hea	alth											
<120>	Diag	nost:	ic ar	nd th	ieraj	eut:	ic ag	geńt:	5						
<130>	2557	460/1	EJH												
<140> <141>	not 2			lable	•		•							•	
<160>	39														
<170>	Pate	ntịn	ver	sion	3.1								•		
<210><211><212><213>	1 1881 DNA human	α	•												<del></del>
<220> <221> <222> <223>	CDS (94)	(1	323)				•								
<400>	1													•	
ataaga	gagg (	ccat	ctgad	ca go	etcca	agata	a cga	acag	cac	tgt	eteca	ata 🤉	gcaad	gatgc	60
Ctaccca	actc (	catca	aagad	ca ga	aaco	ccago	C Ca	Mei 1	g get : Ala	t tog a Sei	g cto	g tgg 1 Trj 5	g gaa o Glu	a tcc 1 Ser	114
Pro Gl	g cag n Gln 10	tgt Cys	atc Ile	atc Ile	ctg Leu	agc Ser 15	cca Pro	ctg Leu	agc Ser	61Å 888	tgg Trp 20	tgg Trp	ttt Phe	tcg Ser	162
atc gga Ile Gly 25	a atc / Ile	tca Ser	ata Ile	ctg Leu	acc Thr 30	agt Ser	tca Ser	gct Ala	ctg Leu	gtg Val 35	ctc Leu	aag Lys	ccc Pro	caa Gln	210
atg cte Met Let 40	aaa Lys	ggc ggc	gaa Glu	ctc Leu 45	cag Gln	act Thr	cga Arg	cct Pro	tct Ser 50	cag Gln	aga Arg	cct Pro	tca Ser	agg Arg 55	258
aag gc	g ttc a Phe	agg Arg	agg Arg 60	aac Asn	aac Asn	ttt Phe	gaa Glu	tat Tyr 65	acc Thr	cta Leu	gaa Glu	gct Ala	tca Ser 70	Lys	306
tca cti Ser Ļei	t cga ı Arg	cag Gln 75	aag Lys	cca Pro	gga Gly	gac Asp	agt Ser 80	acc Thr	atg Met	acg Thr	tac Tyr	ctg Leu 85	aac Asn	aaa Lys	354
ggc cas	ttc phe 90	tat Tyr	ccc Pro	atc Ile	acc Thr	ttg Leu 95	aag Lys	gag Glu	gtg Val	agc Ser	agc Ser 100	agt Ser	gaa Glu	gga Gly	402

Ile	cat His 105	cat His	ccc Pro	atc Ile	agc Ser	aaa Lys 110	gtt Val	cga Arg	agt Ser	gtg Val	atc Ile 115	atg Met	gtg Val	gtt Val	ttt Phe	450
gct Ala 120	gaa Glu	gac Asp	aaa Lys	agc Ser	aga Arg 125	gaa Glu	gat Asp	cag Gln	tta Leu	agg Arg 130	cat His	tgg Trp	aag Lys	tac Tyr	tgg Trp 135	498
cac His	tcc Ser	cgg Arg	cag Gln	cac His	acc Thr	gct Ala	aaa Lys	caa Gln	aga Arg 145	tgc Cys	att Ile	gac Asp	ata Ile	gct Ala 150	gac Asp	546
tat Tyr	aaa Lys	gaa Glu	ago Ser 155	Phe	aac Asp	act Thr	atc Ile	agt Ser 160	Asn	atc Ile	gag Glu	gag Glu	att Ile 165	gcg Ala	tat Tyr	594
aac Asn	gcc Ala	att Ile 170	e Se	tto Phe	aca Thr	tgg Trp	gao Asp 175	Ile	aac Asn	gat Asp	gaa Glu	gca Ala 180	Lys	gtt Val	ttc Phe	642
atc Ile	tct Ser 185	Va.	g aad L Asi	tge n Cy	c tta s Lev	a ago 1 Sei 190	Thr	gat Asp	tto Phe	tct Ser	tcc Ser 195	Glr	g aag 1 Lys	gga gga	gtg Val	690
aaģ Lys	Gly	tt: Le	g cc u Pr	t ct o Le	t aad u Asi 20!	n Ile	c caa	a gtt n Val	gat L Asj	acc Thi 210	Ty:	agt r Sei	t tac r Tyl	aac Asr	aac Asn 215	738
ego Arç	ago g Sei	c aa c As	c aa n Ly	g cc s Pr 22	o Va	g ca l Hi	s Ar	g gc	ta a Ty: 22	r Cy	c cag s Gl:	g aton	c aag e Lys	g gto s Val 230	ttc l Phe	786
tgt Cyt	ga As	c aa p Ly	g gg G G1 23	y Al	t ga a Gl	g cg u Ar	g aa g Ly	a ates Il	e Ar	g ga	t ga p Gl	a ga u Gl	a cg u Ar 24	g Ly	g caa s Gln	834
age Se:	c aa r Ly	a ag s Ar 25	g Ly	a gt 78 Va	t to 1 Se	t ga r As	t gt p Va 25	l Ly	a gt s Va	g cc l Pr	a ct o Le	g ct u Le 26	u Pr	c'tc o Se	t cac r His	882
aa Ly	g cg s Ar 26	g Me	g ga et A	at at	c ac le Th	a gt ir Va 27	l Ph	c aa le Ly	a co 's Pr	c tt	c at e Il 27	e As	it ct sp Le	c ga	t act p Thr	930
ca G1 28	n Pi	t gi	al L	tc t eu P	he I	t co le Pi 85	et ga ro As	ic gt sp Va	g ca al Hi	ac tt Ls Ph 29	ne Al	ec aa La As	ac tt sn Le	g ca u Gl	ng cgg n Arg 295	3
G1 99	c ac y Tì	et c ir H	at g is V	al L	tt co eu P: 00	cc at	tt go le A	cc to la Se	er G	aa ga lu Gi 05	aa ti lu L	tg ga eu G	ag gg lu G	Ly G	aa ggo lu Gly lo	2 1026 Y
to Se	et g	tc t al L	eu I	aa a ye A	gg g rg G	gg c ly P	cg to	yr G	gc a ly T 20	ca g hr G	aa g lu A	at g sp A	sp P	tt ge he Al 25	ct gto la Vai	c 1074 1

cct cct tct acc aag ctg gcc cgg ata gaa gaa cca aag aga gtg ctg Pro Pro Ser Thr Lys Leu Ala Arg Ile Glu Glu Pro Lys Arg Val Leu 330 . 335 340	1122
ctc tac gtt cga aag gag tca gaa gaa gtc ttt gat gcc ctg atg ctc Leu Tyr Val Arg Lys Glu Ser Glu Glu Val Phe Asp Ala Leu Met Leu 345 350 355	1170
aaa acc cca tct ttg aag ggc ttg atg gaa gct atc tca gac aaa tac Lys Thr Pro Ser Leu Lys Gly Leu Met Glu Ala Ile Ser Asp Lys Tyr 360 375	1218
gat gtt ccc cat gac aag att ggg aaa ata ttc aag aag tgt aaa aag Asp Val Pro His Asp Lys Ile Gly Lys Ile Phe Lys Lys Cys Lys Lys 380: 385 390	1266
ggg atc ctg gtg aac atg gac gac aac att gtg aag cat tac tcc aat Gly Ile Leu Val Asn Met Asp Asp Asn Ile Val Lys His Tyr Ser Asn 395 400 405	1314
gag gac acc ttccagctgc agattgaaga agccggggggg tcttacaagc Glu Asp Thr 410	<b>1363</b>
tcaccetgae ggagatetaa aggeetgegg gecaeagete eeeaggagtt eagtgeaggt	1423
gtttctagat cttacggttt ggcaactgca ggtaacccca gtcagccatg tcgccagcac	1483
aggtctatgt cgagggaatg ggttccttgc aggttggagg cggggctgca tctggcttgg	1543
tggtagcatt taatctattg cattggtgtt tttcagatga aagagaaatc catataccat	1603
tatgtttgaa tttcctgata tatacaggat ttaaagtgaa aactttattc caagagttaa	1663
cagagtetet gggaagettt aggacatetg etaegttatt tateaaaata ttgggatete	1723
tgccttgtgc ctacagtgtc gtgggcctgc tcgctagcag aagtcagaaa aggcgatagg	1783
cttggcttta aggatttcgt gcccttgcct gaattcagta caactccact gcctcacgtt	1843
agcgggagcg cacctgaaga gtacgggggg agccctct	1881
<210> 2 <211> 410 <212> PRT <213> human	
<400> 2 Met Ala Ser Leu Trp Glu Ser Pro Gln Gln Cys Ile Ile Leu Ser Pro 1 5 10 15	
Leu Ser Gly Trp Trp Phe Ser Ile Gly Ile Ser Ile Leu Thr Ser Ser 20 25 30	
Ala Leu Val Leu Lys Pro Gln Met Leu Lys Gly Glu Leu Gln Thr Arg 35 40 45	

Pro Ser Gln Arg Pro Ser Arg Lys Ala Phe Arg Arg Asn Asn Phe Glu 50 55 60

Tyr Thr Leu Glu Ala Ser Lys Ser Leu Arg Gln Lys Pro Gly Asp Ser 65 70 75 80

Thr Met Thr Tyr Leu Asn Lys Gly Gln Phe Tyr Pro Ile Thr Leu Lys 85 90 95

Glu Val Ser Ser Ser Glu Gly Ile His His Pro Ile Ser Lys Val Arg 100 105 110

Ser Val Ile Met Val Val Phe Ala Glu Asp Lys Ser Arg Glu Asp Gln 115 120 125

Leu Arg His Trp Lys Tyr Trp His Ser Arg Gln His Thr Ala Lys Gln 130 135 140

Arg Cys Ile Asp Ile Ala Asp Tyr Lys Glu Ser Phe Asn Thr Ile Ser 145 150 155 160

Asn Ile Glu Glu Ile Ala Tyr Asn Ala Ile Ser Phe Thr Trp Asp Ile 165 170 175

Asn Asp Glu Ala Lys Val Phe Ile Ser Val Asn Cys Leu Ser Thr Asp 180 185 190

Phe Ser Ser Gln Lys Gly Val Lys Gly Leu Pro Leu Asn Ile Gln Val

Asp Thr Tyr Ser Tyr Asn Asn Arg Ser Asn Lys Pro Val His Arg Ala 210 215 220

Tyr Cys Gln Ile Lys Val Phe Cys Asp Lys Gly Ala Glu Arg Lys Ile 225 230 235 240

Arg Asp Glu Glu Arg Lys Gln Ser Lys Arg Lys Val Ser Asp Val Lys 245 250 255

Val Pro Leu Leu Pro Ser His Lys Arg Met Asp Ile Thr Val Phe Lys 260 265 270

Pro Phe Ile Asp Leu Asp Thr Gln Pro Val Leu Phe Ile Pro Asp Val 275 280 285

His Phe Ala Asn Leu Gln Arg Gly Thr His Val Leu Pro Ile Ala Ser 290 295 300

Glu Glu Leu Glu Gly Glu Gly Ser Val Leu Lys Arg Gly Pro Tyr Gly 305 310 315 . 320

Thr Glu Asp Asp Phe Ala Val Pro Pro Ser Thr Lys Leu Ala Arg Ile 325 330 335

Glu Glu Pro Lys Arg Val Leu Leu Tyr Val Arg Lys Glu Ser Glu Glu

			340					345					350			•
Val	Phe	Asp 355	Ala	Leu	Met	Leu	<b>Ъу</b> в 360	Thr	Pro	Ser	Leu	<b>L</b> ys 365	Gly	Leu	Met	
Glu	Ala 370	Ile	Ser	Asp	Lys	Tyr 375	qaA	Val	Pro	His	Asp 380	Lув	Ile	Gly	Lys	
Ile 385	Phe	Lys	Lys	Сув	390	ГÀЗ	Gly	Ile	Leu	Val 395	Asn	Met	Asp	Asp	Asn 400	
Ile	Val	Lys	His	Tyr 405	Ser	Asn	Glu	Asp	Thr 410	•						
<210 <211 <212 <213	l> 2 2> 1	3 2361 ONA numar	1										-			
<220 <221 <222 <223	L> ( 2>	CDS (7)	. (186	50)								٠	•			
<400 agc	gog a	3 atg <i>a</i> Met 1	aca c	ag g	gag t 3lu 7	ac g	ac a	ac a	aaa d Lys 2	egg (	cca (	gtg ( Val )	Ltg (	gtt d Val 1	ett Leu	48
		1				5					10					
cag Gln 15	aat	gaa Glu	gca	ctt	tat	cca	caq	caa	caa	tee:	tac	act	aat	asa	ast	96
15 gag	aat Asn	gaa	gca Ala aaa	ctt Leu tcc	tat Tyr 20	cca Pro	cag Gln gaa	cgg Arg	cgg Arg	tcc Ser 25	tac Tyr	act Thr	agt Ser	gag Glu	gat Asp 30	96 144
gag Glu gcg	aat Asn gcc Ala	gaa Glu tgg	gca Ala aaa Lys	ctt Leu tcc Ser 35	tat Tyr 20 ttc Phe	cca Pro ctg Leu	cag Gln gaa Glu	cgg Arg aac Asn	cgg Arg cct Pro 40	tcc Ser 25 ctc Leu	tac Tyr act Thr	act Thr gca Ala	agt Ser gcg Ala	gag Glu acc Thr 45	gat Asp 30 aaa Lys	
gag Glu gcg Ala	aat Asn gcc Ala atg Met	gaa Glu tgg Trp	gca Ala aaa Lys agc Ser 50	ctt Leu tcc Ser 35 atc Ile	tat Tyr 20 ttc Phe aat Asn	cca Pro ctg Leu gga Gly	cag Gln gaa Glu gat Asp	cgg Arg aac Asn gaa Glu 55	cgg Arg cct Pro 40 gac Asp	tcc Ser 25 ctc Leu	tac Tyr act Thr gcc Ala	act Thr gca Ala gct Ala	agt Ser gcg Ala gcg Ala 60	gag Glu acc Thr 45 ctg Leu	gat Asp 30 aaa Lys ggc Gly	144
gag Glu gcg Ala ctg Leu	aat Asn gcc Ala atg Met ctc Leu	gaa Glu tgg Trp atg Met tat	gca Ala aaa Lys agc Ser 50 gac Asp	ctt Leu tcc Ser 35 atc Ile tac Tyr	tat Tyr 20 ttc Phe aat Asn tac Tyr	cca Pro ctg Leu gga Gly aag Lys	cag Gln gaa Glu gat Asp gtt Val 70	cgg Arg aac Asn gaa Glu 55 cca Pro	cgg Arg cct Pro 40 gac Asp aga Arg	tcc ser 25 ctc Leu agc ser gag Glu	tac Tyr act Thr gcc Ala aga Arg	act Thr gca Ala gct Ala agg Arg 75	agt Ser gcg Ala gcg Ala 60 tca	gag Glu acc Thr 45 ctg Leu tca ser	gat Asp 30 aaa Lys ggc Gly aca Thr	144 192
gag Glu gcg Ala ctg Leu gca Ala	aat Asn gcc Ala atg Met ctc Leu aag Lys 80 ata	gaa Glu tgg Trp atg Met tat Tyr 65 cca	gca Ala aaa Lys agc Ser 50 gac Asp	ctt Leu tcc Ser 35 atc Ile tac Tyr	tat Tyr 20 ttc Phe aat Asn tac Tyr	cca Pro ctg Leu gga Gly aag Lys cac His	cag Gln gaa Glu gat Asp gtt Val 70 cct Pro	cgg Arg aac Asn gaa Glu 55 cca Pro	cgg Arg cct Pro 40 gac Asp aga Arg	tcc ser 25 ctc Leu agc ser gag Glu gat Asp	tac Tyr act Thr gcc Ala aga Arg cac His	act Thr gca Ala gct Ala agg Arg 75 agc Ser	agt Ser gcg Ala gcg Ala 60 tca ser aaa Lys	gag Glu acc Thr 45 ctg Leu tca ser aga Arg	gat Asp 30 aaa Lys ggc Gly aca Thr	144 192 240

Gly	aac Asn	cag Gln	ctg Leu 130	ggc	att Ile	gat Asp	aag Lys	aga Arg 135	ggc Gly	cat His	ctg Leu	aca Thr	gct Ala 140	tca Ser	gat Asp	432
acg Thr	aca Thr	gtc Val 145	act Thr	gtc Val	tcc Ser	ata Ile	gca Ala 150	acg Thr	atg Met	cct Pro	acc Thr	cac His 155	Ser	atc Ile	aag Lys	480
aca Thr	gaa Glu 160	acc Thr	cag Gln	cca Pro	cat His	ggc Gly 165	ttc Phe	gct Ala	gtg Val	gga Gly	atc Ile 170	ccc Pro	cca Pro	gca Ala	gtg Val	528
tat Tyr 175	cat His	cct Pro	gag Glu	ccc Pro	act Thr 180	gag Glu	cgg Arg	gtg Val	gtg Val	gtt Val 185	ttc Phe	gat Asp	egg Cgg	aay Asn	ctc Leu 190	576
aat Asn	act Thr	gac Asp	cag Gln	ttc Phe 195	agc Ser	tct Ser	ggt	gct Ala	caa Gln 200	gcc Ala	cca Pro	aat Asn	gct Ala	caa Gln 205	agg Arg	624
cga Arg	act Thr	cca Pro	gac Asp 210	tcg Ser	acc Thr	ttc Phe	tca Ser	gag Glu 215	acc Thr	ttc Phe	aag Lys	gaa Glu	ggc Gly 220	gtt Val	cag Gln	672
gag Glu	gtt Val	ttc Phe 225	ttc Phe	ccc Pro	tcg Ser	gat Asp	ctc Leu 230	agt Ser	ctg Leu	cgg Arg	atg Met	cct Pro 235	ggc Gly	atg Met	aat Asn	720
tca Ser	gag Glu 240	gac Asp	tat Tyr	gtt Val	ttt Phe	gac Asp 245	agt Ser	gtt Val	tct Ser	ejå aaa	aac Asn 250	aac Asn	ttt Phe	gaa Glu	tat Tyr	768
acc Thr 255	cta Leu	gaa Glu	gct Ala	tca Ser	aaa Lys 260	tca Ser	ctt Leu	cga Arg	cag Gln	aag Lys 265	cca Pro	gga Gly	gac Asp	agt Ser	acc Thr 270	816
atg Met	acg Thr	tac Tyr	ctg Leu	aac Asn 275	aaa Lys	ggc Gly	cag Gln	ttc Phe	tat Tyr 280	ccc Pro	atc Ile	acc Thr	ttg Leu	aag Lys 285	gag Glu	864
gtg Val	agc Ser	agc Ser	agt Ser 290	gaa Glu	gga Gly	atc Ile	cat His	cat His 295	ccc Pro	atc Ile	agc Ser	aaa Lys	gtt Val 300	cga Arg	agt Ser	912
gtg Val	atc Ile	atg Met 305	gtg Val	gtt Val	ttt Phe	gct Ala	gaa Glu 310	gac Asp	aaa Lys	agc Ser	aga Arg	gaa Glu 315	gat Asp	cag Gln	tta Leu	960
agg Arg	cat His 320	tgg Trp	aag Lys	tac Tyr	tgg Trp	cac His 325	tcc Ser	cgg Arg	cag Gln	cac His	acc Thr 330	gct Ala	aaa Lys	caa Gln	aga Arg	1008
tgc Cys 335	att Ile	gac Asp	ata Ile	gct Ala	gac Asp 340	tat Tyr	awa Xaa	gaa Glu	agc Ser	ttc Phe 345	aac Asn	act Thr	atc Ile	agt Ser	aac Asn 350	1056

atc Ile	gag Glu	gag Glu	att Ile	gcg Ala 355	tat Tyr	aac Asn	gcc Ala	att Ile	tcc Ser 360	ttc Phe	aca Thr	tgg Trp	gac Asp	atc Ile 365	aac Asn	1104
gat Asp	gaa Glu	gca Ala	aag Lys 370	gtt Val	ttc Phe	atc Ile	tct Ser	gtg Val 375	aac Asn	tgc Cys	tta Leu	agc Ser	aca Thr 380	gat Asp	ttc Phe	1152
tct Ser	tcc Ser	cag Gln 385	aag Lys	gga Gly	gtg Val	aag Lys	999 Gly 390	ttg Leu	cct Pro	ctt Leu	aac Asn	att Ile 395	caa Gln	gtt Val	gat Asp	1200
acc Thr	tat Tyr 400	agt Ser	tac Tyr	aac Asn	aac Asn	cgc Arg 405	agc Ser	aac Asn	aag Lys	cct Pro	gtg Val 410	cac His	cgg Arg	gcc Ala	tac Tyr	1248
tgc Cys 415	cag Gln	atc Ile	aag Lys	gtc Val	ttc Phe 420	tgt Cys	gac Asp	aag Lys	gga Gly	gct Ala 425	gag Glu	cgg	aaa Lys	atc Ile	agg Arg 430	1296
Asp	Glu	Glu	Arg	Lys 435	Gln	Ser	Lys	Arg	Lys 440	Val	Ser	gat Asp	Val	Lys 445	Val	1344
Pro	Leu	Leu	Pro 450	Ser ··	His	ГÀв	Arg	Met 455	qaA	Ile	Thr	gtt Val	Phe 460	Lys	Pro	1392
Phe	Ile	Asp 465	Leu	Asp	Thr	Gln	Pro 470	Val	Leu	Phe	,Ile	Pro 475	Āsp	Val	His	1440
Phe	Ala 480	Asn	Leu	Gln	Arg	Gly 485	Thr	His	.Val	Leu	490	lle	Ala	Ser	gaa Glu	1488
Glu 495	Leu	Glu	Gly	Glu	Gly 500	Ser	. Val	. Leu	Lys	Arg 505	Gly	Pro	Тух	Gly	Thr 510	1536
Glu	Asp	Asp	Phe	515	Val	Pro	Pro	Sei	Thr 520	Lys	Let	a Ala	Arg	525		1584
Glu	Pro	Lys	530	y Va]	. Lev	Lev	ι Туз	7 Val	L Arg	l Lys	Gli	ı Sez	540	Glu	y gtc	1632
Phe	e Asp	9 Ala 549	a Let	ı Met	: Lev	ı Lya	550	r Pro	Se:	Lev	і Ьу:	55!	Let	ı Met	g gaa : Glu	1680
Ala	a Ile	e Se	a gad r Ası	aaa Lys	a tac s Tyr	gat Asp 569	va:	l Pro	c cat	z gad s Asp	Ly 57	s Ile	• Gl	g aaa / Lys	a ata 3 Ile	1728

ttc aag aag tgt-aaa aag ggg atc ctg gtg aac atg gac gac aac att Phe Lys Lys Cys Lys Gly Ile Leu Val Asn Met Asp Asp Asn Ile 575 580 585	1776
gtg aag cat tac tcc aat gag gac acc ttc cag ctg cag att gaa gaa Val Lys His Tyr Ser Asn Glu Asp Thr Phe Gln Leu Gln Ile Glu Glu 595 600 605	1824
gcc ggg ggg tct tac aag ctc acc ctg acg gag atc taaaggcctg Ala Gly Gly Ser Tyr Lys Leu Thr Leu Thr Glu Ile 610 615	1870
cgggccacag ctccccagga gttcagtgca ggtgtttcta.gatcttacgg tttggcaact	1930
gcaggtaacc ccagtcagcc atgtcgccag cacaggtcta tgtcgaggga atgggttcct	1990
tgcaggttgg aggcggggct gcatctggct tggtggtagc atttaatcta ttgcattggt	2050
gtttttcaga tgaaagagaa atccatatac cattatgttt gaatttcctg atatatacag	2110
gatttaaagt gaaaacttta ttccaagagt taacagagtc tctgggaagc tttaggacat	2170
ctgctacgtt atttatcaaa atattgggat ctctgccttg tgcctacagt gtcgtgggcc	2230
tgetegetag cagaagteag aaaaggegat aggettgget ttaaggattt egtgeeettg	2290
cetgaattea gtacaaetee aetgeeteae gttageggga gegeaeetga agagtaeggg	2350
gggagecete t	2361
<210> 4 <211> 618 <212> PRT <213> human	
<220> <221> misc_feature <222> (342)(342) <223> The 'Xaa' at location 342 stands for Lys, or Ile.	
<pre>&lt;400&gt; 4 Met Thr Gln Glu Tyr Asp Asn Lys Arg Pro Val Leu Val Leu Gln Asn 1 5 10 15</pre>	
Glu Ala Leu Tyr Pro Gln Arg Arg Ser Tyr Thr Ser Glu Asp Glu Ala 20 25 30	
Trp Lys Ser Phe Leu Glu Asn Pro Leu Thr Ala Ala Thr Lys Ala Met 35 40 45	
Met Ser Ile Asn Gly Asp Glu Asp Ser Ala Ala Ala Leu Gly Leu Leu 50 55 60	
Tyr Asp Tyr Tyr Lys Val Pro Arg Glu Arg Arg Ser Ser Thr Ala Lys 65 70 . 75 80	

Pro Glu Val Glu His Pro Glu Pro Asp His Ser Lys Arg Asn Ser Ile Pro Ile Val Thr Glu Gln Pro Leu Ile Ser Ala Gly Glu Asn Arg Val-105 Gln Val Leu Lys Asn Val Pro Phe Asn Ile Val Leu Pro His Gly Asn 120 Gln Leu Gly Ile Asp Lys Arg Gly His Leu Thr Ala Ser Asp Thr Thr Val Thr Val Ser Ile Ala Thr Met Pro Thr His Ser Ile Lys Thr Glu Thr Gln Pro His Gly Phe Ala Val Gly Ile Pro Pro Ala Val Tyr His Pro Glu Pro Thr Glu Arg Val Val Val Phe Asp Arg Asn Leu Asn Thr 180 Asp Gln Phe Ser Ser Gly Ala Gln Ala Pro Asn Ala Gln Arg Arg Thr 200 Pro Asp Ser Thr Phe Ser Glu Thr Phe Lys Glu Gly Val Gln Glu Val Phe Phe Pro Ser Asp Leu Ser Leu Arg Met Pro Gly Met Asn Ser Glu 230 225 . Asp Tyr Val Phe Asp Ser Val Ser Gly Asn Asn Phe Glu Tyr Thr Leu 250 Glu Ala Ser Lys Ser Leu Arg Gln Lys Pro Gly Asp Ser Thr Met Thr Tyr Leu Asn Lys Gly Gln Phe Tyr Pro Ile Thr Leu Lys Glu Val Ser Ser Ser Glu Gly Ile His His Pro Ile Ser Lys Val Arg Ser Val Ile Met Val Val Phe Ala Glu Asp Lys Ser Arg Glu Asp Gln Leu Arg His 315 Trp Lys Tyr Trp His Ser Arg Gln His Thr Ala Lys Gln Arg Cys Ile Asp Ile Ala Asp Tyr Xaa Glu Ser Phe Asn Thr Ile Ser Asn Ile Glu 345 Glu Ile Ala Tyr Asn Ala Ile Ser Phe Thr Trp Asp Ile Asn Asp Glu 360

Ala Lys Val Phe Ile Ser Val Asn Cys Leu Ser Thr Asp Phe Ser Ser

<220><221>

<222>

<223>

CDS

(67) . . (1941)

370 375 380 Gln Lys Gly Val Lys Gly Leu Pro Leu Asn Ile Gln Val Asp Thr Tyr 390 395 Ser Tyr Asn Asn Arg Ser Asn Lys Pro Val His Arg Ala Tyr Cys Gln Ile Lys Val Phe Cys Asp Lys Gly Ala Glu Arg Lys Ile Arg Asp Glu Glu Arg Lys Gln Ser Lys Arg Lys Val Ser Asp Val Lys Val Pro Leu 440 Leu Pro Ser His Lys Arg Met Asp Ile Thr Val Phe Lys Pro Phe Ile Asp Leu Asp Thr Gln Pro Val Leu Phe Ile Pro Asp Val His Phe Ala Asn Leu Gln Arg Gly Thr His Val Leu Pro Ile Ala Ser Glu Glu Leu 485 Glu Gly Glu Gly Ser Val Leu Lys Arg Gly Pro Tyr Gly Thr Glu Asp Asp Phe Ala Val Pro Pro Ser Thr Lys Leu Ala Arg Ile Glu Glu Pro Lys Arg Val Leu Leu Tyr Val Arg Lys Glu Ser Glu Glu Val Phe Asp 535 Ala Leu Met Leu Lys Thr Pro Ser Leu Lys Gly Leu Met Glu Ala Ile Ser Asp Lys Tyr Asp Val Pro His Asp Lys Ile Gly Lys Ile Phe Lys Lys Cys Lys Lys Gly Ile Leu Val Asn Met Asp Asp Asn Ile Val Lys 585 His Tyr Ser Asn Glu Asp Thr Phe Gln Leu Gln Ile Glu Glu Ala Gly Gly Ser Tyr Lys Leu Thr Leu Thr Glu Ile 615 <210> 5 <211> 4532 <212> DNA <213> human

<400> 5 ttgaaagtcc agtttcacca gaggctgagg ctccaggaaa aggggagcaa gttcattgga	60
tcaaac atg tca caa gag tca gac aat aat aaa aga cta gtg gcc tta Met Ser Gln Glu Ser Asp Asn Asn Lys Arg Leu Val Ala Leu 1 5 10	108
gtg ccc atg ccc agt gac cct cca ttc aat acc cga aga gcc tac acc Val Pro Met Pro Ser Asp Pro Pro Phe Asn Thr Arg Arg Ala Tyr Thr 15 20 25 30	<b>156</b>
agt gag gat gaa gcc tgg aag tca tac ttg gag aat ccc ctg aca gca Ser Glu Asp Glu Ala Trp Lys Ser Tyr Leu Glu Asn Pro Leu Thr Ala 35 40 45	204
gcc acc aag gcc atg atg agc att aat ggt gat gag gac agt gct Ala Thr Lys Ala Met Met Ser Ile Asn Gly Asp Glu Asp Ser Ala Ala 50 55 60	252
gcc ctc ggc ctg ctc tat gac tac tac aag gtt cct cga gac aag agg Ala Leu Gly Leu Leu Tyr Asp Tyr Tyr Lys Val Pro Arg Asp Lys Arg 65 70 75	300
ctg ctg tct gta agc aaa gca agt gac agc caa gaa gac cag gag aaa Leu Leu Ser Val Ser Lys Ala Ser Asp Ser Gln Glu Asp Gln Glu Lys 80 85 90	348
aga aac tgc ctt ggc acc agt gaa gcc cag agt aat ttg agt gga gga Arg Asn Cys Leu Gly Thr Ser Glu Ala Gln Ser Asn Leu Ser Gly Gly 95 100 105 110	396
gaa aac cga gtg caa gtc cta aag act gtt cca gtg aac ctt tcc cta Glu Asn Arg Val Gln Val Leu Lys Thr Val Pro Val Asn Leu Ser Leu 115 120 125	444
aat caa gat cac ctg gag aat tcc aag cgg gaa cag tac agc atc agc Asn Gln Asp His Leu Glu Asn Ser Lys Arg Glu Gln Tyr Ser Ile Ser 130 135	492
ttc ccc gag agc tct gcc atc atc ccg gtg tcg gga atc acg gtg gtg Phe Pro Glu Ser Ser Ala Ile Ile Pro Val Ser Gly Ile Thr Val Val 145 150 155	540
aaa gct gaa gat ttc aca cca gtt ttc atg gcc cca cct gtg cac tat Lys Ala Glu Asp Phe Thr Pro Val Phe Met Ala Pro Pro Val His Tyr 160 165 170	588
ccc cgg gga gat ggg gaa gag caa cga gtg gtt atc ttt gaa cag act Pro Arg Gly Asp Gly Glu Glu Gln Arg Val Val Ile Phe Glu Gln Thr 175 180 185 190	636
cag tat gac gtg ccc tcg ctg gcc acc cac agc gcc tat ctc aaa gac Gln Tyr Asp Val Pro Ser Leu Ala Thr His Ser Ala Tyr Leu Lys Asp 195 200 205	684

gac Asp	cag Gln	cgc Arg	agc Ser 210	act Thr	ccg Pro	gac Asp	agc Ser	aca Thr 215	tac Tyr	agc Ser	gag Glu	agc Ser	ttc Phe 220	aag Lys	gac Asp	732
gca Ala	gcc Ala	aca Thr 225	gag Glu	aaa Lys	ttt Phe	cgg Arg	agt Ser 230	gct Ala	tca Ser	gtt Val	GJA aaa	gct Ala 235	gag Glu	gag Glu	tac Tyr	780
atg Met	tat Tyr 240	gat Asp	cag Gln	aca Thr	tca Ser	agt Ser 245	Gly	aca Thr	ttt Phe	cag Gln	tac Tyr 250	acc Thr	ctg Leu	gaa Glu	gcc Ala	828
acc Thr 255	aaa Lys	tct Ser	ctc Leu	cgt Arg	cag Gln 260	aag Lys	cag Gln	Gly 999	gag Glu	ggc Gly 265	ccc Pro	atg Met	acc Thr	tac Tyr	ctc Leu 270	876
Asn	Lys	Gly	Gln	Phe 275	Tyr	Ala	Ile	aca Thr	Leu 280	Ser	Glu	Thr	Gly	Asp 285	Asn	924
гàв	Cys	Phe	Arg 290	His	Pro	Ile	Ser	aaa Lys 295	Val	Arg	Ser	Val	Val 300	Met	Val	<b>972</b>
val	Phe	305	Glu	Asp	ГÀв	Asn	Arg 310	gat Asp	Glu	Gln	Leu	Lys 315	Tyr	Trp	Lys	1020
Tyr	320	His	Ser	Arg	Gln	His 325	Thr	gcg Ala	Lys	Gln	Arg 330	Val	Leu	Asp	Ile	1068
335	Asp	Tyr	ГÀв	Glu	Ser 340	Phe	Asn	acg Thr	Ile	Gly 345	Asn	Ile	Glu	Glu	Ile 350	1116
ATa	Tyr	Asn	Ala	Val 355	Ser	Phe	Thr	tgg Trp	360	Val	Asn	Glu	Glu	Ala 365	Lys	1164
тте	Phe	Ile	370	Val	Asn	Сув	Leu	agc Ser 375	Thr	Asp	Phe	Ser	Ser 380	Gln	Lys	1212
GIĀ	Val	195 385	GIÀ	Leu	Pro	Leu	Met 390		Gln	Ile	Asp	Thr 395	Tyr	Ser	Tyr	1260
Asn	Asn 400	Arg	Ser	Asn	Lys	Pro 405	Ile	His	Arg	Ala	Tyr 410	Сув	Gln	Ile	aag Lys	1308
gtc Val 415	hue	tgt Cys	gac Asp	aaa Lys	gga Gly 420	gca Ala	gaa Glu	aga Arg	aaa Lys	atc Ile 425	Arg	gat Asp	gaa Glu	gag Glu	cgg Arg 430	1356

aag Lys	cag Gln	aac Asn	agg Arg	aag Lys 435	aaa Lys	Gly 999	aaa Lys	ggc	cag Gln 440	gcc Ala	tcc Ser	caa Gln	act Thr	caa Gln 445	tgc Cys	1404
aac Asn	agc Ser	tcc Ser	tct Ser 450	gat Asp	Gly aaa	aag Lys	ttg Leu	gct Ala 455	gcc Ala	ata <sup>.</sup> Ile	cct Pro	tta Leu	cag Gln 460	aag Lys	aag Lys	1452
agt Ser	gac Asp	atc Ile 465	acc Thr	tac Tyr	ttc Phe	aaa Lys	acc Thr 470	atg Met	cct Pro	gat Asp	ctc Leu	cac His 475	tca Ser	cag Gln	cca Pro	1500
gtt Val	ctc Leu 480	Pue	ata Ile	cct Pro	gat Asp	gtt Val 485	cac His	ttt Phe	gca Ala	aac Asn	ctg Leu 490	cag Gln	agg Arg	acc Thr	gga Gly	1548
cag Gln 495	gtg Val	tat Tyr	tac Tyr	aac Asn	acg Thr 500	gat Asp	gat Asp	gaa Glu	cga Arg	gaa Glu 505	ggt Gly	ggc Gly	agt Ser	gtc Val	ctt Leu 510	1596
gtt Val	aaa Lys	cgg Arg	atg Met	ttc Phe 515	cgg Arg	ccc Pro	atg Met	gaa Glu	gag Glu 520	gag Glu	ttt Phe	ggt Gly	cca Pro	gtg Val 525	cct Pro	1644
SCI	пув	GIN	530	гув	GIU	GIu	Gly	aca Thr 535	Lys	Arg	Val	Leu	Leu 540	Tyr	Val	1692
agg Arg	aag Lys	gag Glu 545	act Thr	gac Asp	gat Asp	gtg Val	ttc Phe 550	gat Asp	gca Ala	ttg Leu	atg Met	ttg Leu 555	aag Lys	tct Ser	ccc Pro	1740
aca Thr	gtg Val 560	aag Lys	Gly	ctg Leu	Met	gaa Glu 565	gcg Ala	ata Ile	tct Ser	gag Glu	aaa Lys 570	tat Tyr	gly ggg	ctg Leu	ccc Pro	1788
gtg Val 575	gag Glu	aag Lys	ata Ile	gca Ala	aag Lys 580	ctt Leu	tac Tyr	aag Lys	aaa Lys	agc Ser 585	aaa Lys	aaa Lys	ggc Gly	atc Ile	ttg ' Leu 590	1836
gtg Val	aac Asn	atg Met	gat Asp	gac Asp 595	aac Asn	atc Ile	atc Ile	gag Glu	cac His 600	tac Tyr	tcg Ser	aac Asn	gag Glu	gac Asp 605	acc Thr	1884
ttc Phe	atc Ile	ctc Leu	aac Asn 610	atg Met	gag Glu	agc Ser	atg Met	gtg Val 615	gag Glu	ggc Gly	ttc Phe	aag Lys	gtc Val 620	acg Thr	ctc Leu	1932
atg Met	gaa Glu	atc Ile 625	tage	ccct	ggg (	ttg	gcat	cc g	cttt	ggct	g gag	getei	tcag			1981
tgc	gttc	ete d	cctg	agaga	ag a	caga	agcc	c ca	gccc	caga	acci	tgga	gac o	ccat	ctcccc	2041
cate	ctca	caa d	ctgc	tgtta	ac a	agac	egtg	c tg	ggga	gtgg	ggca	aagg	gac a	aggc	ccact	2101

gteggtgtge ttggcccate caetggcace taccaeggag ctgaagcetg agcccctcag 2161 gaaggtgcct taggcctgtt ggattcctat ttattgccca ccttttcctg gagcccaggt 2221 ccaggcccgc caggactctg caggtcactg ctagctccag atgagaccgt ccagcgttcc 2281 cccttcaaga gaaacactca tcccgaacag cctaaaaaat tcccatccct tctctctcac 2341 2401 ccctccatat ctatctcccg agtggctgga caaaatgagc tacgtctggg tgcagtagtt ataggtgggg caagaggtgg atgcccactt tctggtcaga cacctttagg ttgctctggg 2461 gaaggetgte ttgetaaata ceteeagggt teecageaag tggeeaceag geettgtaca 2521 ggaagacatt cagtcaccgt gtaattagta acacagaaag tetgeetgte tgcattgtac 2581 atagtgttta taatattgta ataatatatt ttacctgtgg tatgtgggca tgtttactgc 2641 cactggcett agaggagaca cagacetgga gacegtttta atgggggttt ttgcctctgt 2701 2761 geetgtteaa gagaettgea gggetaggta gagggeettt gggatgttaa ggtgaetgea 2821 getgatgeca agatggaete tgeaatggge ataeetgggg getegtteee tgteeceaga ggaagcccc teteettete catgggcatg acteteette gaggccacca cgtttatete 2881 acaatgatgt gttttgcttg actttccctt tgcgctgtct cgtgggaaag gtcattctgt 2941 3001 etgagacccc ageteettet ceagetttgg etgegggeat ggeetgaget ttetggagag cetetgeagg gggtttgeea teagggeeet gtggetgggt etgetgeaga geteettgge 3061 3121 tatcaggaga atcctggaca ctgtactgtg cctcccagtt tacaaacacg cccttcatct 3181 caagtggccc tttaaaaggc ctgctgccat gtgagagctg tgaacagctc agctctgagt eggeaggetg gggetteete etgggeeace agatggaaag ggggtattgt ttgeeteact 3241 cctggatgct gcgttttaag gaagtgagtg agaaagaatg tgccaagata cctggctcct 3301 gtgaaaccag cctcaggagg gaaactggga gagagaagct gtggtctcct gctacatgcc 3361 3421 ctgggagctg gaagagaaaa acactcccct aaacaatcgc aaaatgatga accatcatgg 3481 gecactgtte tetttgaggg gacaggttta ggggtttgeg ttegecettg tgggetgaag 3541 cactagettt ttggtageta gacacateet geacecaaag gttetetaea aaggeecaga 3601 tttgtttgta aagcactttg actottacct ggaggcccgc tctctaaggg cttcctgcgc teccaectea tetgteeetg agatgeagag eaggatggag ggtetgette tageteaget 3661 3721 gtttctcctt gaggttgcgg aggaattgaa ttgaatggga cagagggcag gtgctgtggc 3781 caagaagatc teegageage agtgaegggg caeettgetg tgtgteetet gggeatgtta

accettetgt ggggccaaag gtttgcatcg tggatccage tgtgctccag tctgtcccct	3841
cetectecae tetgaetgee aegeecegga ceageagett ggggaecete cagggtaeta	3901
atggggetet gttetgagat ggacaaatte agtgttggaa atacatgttg tactatgcae	3961
ttcccatgct cctagggtta ggaatagttt caaacatgat tggcagacat aacaacggca	4021
aatactcgga ctggggcata ggactccaga gtaggaaaaa gacaaaagat ttggcagcct	4081
gacacaggca acctacccct ctctctccag cctctttatg aaactgtttg tttgccagtc	4141
ctgccctaag gcagaagatg aattgaagat gctgtgcatg tttcctaagt ccttgagcaa	4201
tcatggtggt gacaattgcc acaagggata tgaggccagt gccaccagag ggtggtgcca	4261
agtgccacat cccttccgat ccattcccct ctgcatcctc ggagcacccc agtttgcctt	4321
tgatgtgtcc gctgtgtatg ttagctgaac tttgatgagc aaaatttcct gagcgaaaca	4381
etecaaagag ataggaaaac ttgccgcctc ttcttttttg tcccttaatc aaactcaaat	4441
aagettaaaa aaaateeatg gaagateatg gacatgtgaa atgageattt ttttetttt	4501
ttttttttt tttaacaag tctgaactga g	4532
<210> 6 <211> 625 <212> PRT <213> human	٠.
<400> 6	
Met Ser Gln Glu Ser Asp Asn Asn Lys Arg Leu Val Ala Leu Val Pro  1 10 15	
Met Pro Ser Asp Pro Pro Phe Asn Thr Arg Arg Ala Tyr Thr Ser Glu 20 25 ' 30	٠.
Asp Glu Ala Trp Lys Ser Tyr Leu Glu Asn Pro Leu Thr Ala Ala Thr 35 40 45	
Lys Ala Met Met Ser Ile Asn Gly Asp Glu Asp Ser Ala Ala Leu 50 55 60	•
Gly Leu Leu Tyr Asp Tyr Tyr Lys Val Pro Arg Asp Lys Arg Leu Leu 65 70 75 80	
Ser Val Ser Lys Ala Ser Asp Ser Gln Glu Asp Gln Glu Lys Arg Asn 85 90 95	
Cys Leu Gly Thr Ser Glu Ala Gln Ser Asn Leu Ser Gly Glu Asn 100 105 110	
Arg Val Gln Val Leu Lys Thr Val Pro Val Asn Leu Ser Leu Asn Gln 115 120 125	

120

Asp His Leu Glu Asn Ser Lys Arg Glu Gln Tyr Ser Ile Ser Phe Pro Glu Ser Ser Ala Ile Ile Pro Val Ser Gly Ile Thr Val Val Lys Ala Glu Asp Phe Thr Pro Val Phe Met Ala Pro Pro Val His Tyr Pro Arg 170 Gly Asp Gly Glu Glu Gln Arg Val Val Ile Phe Glu Gln Thr Gln Tyr Asp Val Pro Ser Leu Ala Thr His Ser Ala Tyr Leu Lys Asp Asp Gln Arg Ser Thr Pro Asp Ser Thr Tyr Ser Glu Ser Phe Lys Asp Ala Ala Thr Glu Lys Phe Arg Ser Ala Ser Val Gly Ala Glu Glu Tyr Met Tyr Asp Gln Thr Ser Ser Gly Thr Phe Gln Tyr Thr Leu Glu Ala Thr Lys Ser Leu Arg Gln Lys Gln Gly Glu Gly Pro Met Thr Tyr Leu Asn Lys Gly Gln Phe Tyr Ala Ile Thr Leu Ser Glu Thr Gly Asp Asn Lys Cys Phe Arg His Pro Ile Ser Lys Val Arg Ser Val Val Met Val Val Phe Ser Glu Asp Lys Asn Arg Asp Glu Gln Leu Lys Tyr Trp Lys Tyr Trp His Ser Arg Gln His Thr Ala Lys Gln Arg Val Leu Asp Ile Ala Asp Tyr Lys Glu Ser Phe Asn Thr Ile Gly Asn Ile Glu Glu Ile Ala Tyr Asn Ala Val Ser Phe Thr Trp Asp Val Asn Glu Glu Ala Lys Ile Phe 360 Ile Thr Val Asn Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Val Lys Gly Leu Pro Leu Met Ile Gln Ile Asp Thr Tyr Ser Tyr Asn Asn 395 Arg Ser Asn Lys Pro Ile His Arg Ala Tyr Cys Gln Ile Lys Val Phe Cys Asp Lys Gly Ala Glu Arg Lys Ile Arg Asp Glu Glu Arg Lys Gln

Asn	Arg	Lys 435	ГÀЗ	Gly	Lys	Gly	Gln 440	Ala	Ser	Gln	Thr	Gln 445	Сув	Asn	Ser
Ser	Ser 450	Asp	Gly	Lys	Leu	Ala 455	Ala	Ile	Pro	Leu	Gln 460	Гуs	Lys	Ser	Asp
Ile 465	Thr	Tyr	Phe	Lys	Thr 470	Met	Pro	Asp	Leu	His 475	Ser	Gln	Pro	Val	Leu 480
Phe	Ile	Pro	Asp	Val 485	His	Phe	Ala	Asn	Leu 490	Gln	Arg	Thr	Gly	Gln 495	Val.
Tyr	Tyr	Asn	Thr 500	Asp	Asp	. Glu	Arg	Glu 505	Gly	Gly	Ser	Val	Leu 510	Val	Lys
Arg	Met	Phe 515	Arg	Pro	Met	Glu	Glu 520	Glu	Phe	Gly	Pro	Val 525	Pro	Ser	PAe
Gln	Met 530		Glu	Glu	Gly	Thr 535	Lys	Arg	Val	Leu	Leu 540	Tyr	Val	Arg	Lys
Glu 545		Asp	Asp	Val	Phe 550	_	Ala	Leu	Met	Leu 555	-	Ser	Pro	Thr	Val 560
Lys	Gly	Leu	Met	Glu 565		Ile	Ser	Glu	<b>Ьуз</b> 570	_	.Gly	Leu	Pro	Val 575	
Lys	Ile	Ala	. <b>Т</b> ув 580		Tyr	ГÀЗ	Lys	Ser 585	-	Lys	Gly	Ile	Ьеи 590		Asn
Met	Asp	Asp 595		Ile	: Ile	Glu	His 600	-	Ser	Asr	Glu	Asp 605		Phe	Ile
Lev	Asr 610		Glu	. Ser	Met	Val 615		Gly	Phe	Lys	620		Leu	Met	Glu
Ile 625				•		•									
<23		7													•
	l1> l2>	1870 DNA	,									•			
	13>	huma	an												
<2	20 <i>5</i>														
	21>	_													
	22> 23>	(47)	) (:	1867)	)										
<4	00>	7													
			cca	aact	gtt a	aaga	gtgg	tt a	tttc	tgag	c ag	aaga			
	٠						•			•			Met 1	Trp	Met
aa	t tc	c at	t ct	t cc	t at	t tt	t ct	t tt	c ag	g tc	t gt	g cg	g ct	g cta	a aag

55

Asn	Ser 5	Ile	Leu	Pro	Ile	Phe 10	Leu	Phe	Arg	Ser	Val 15	Arg	Leu	Leu	Lys	
aac Asn 20	gac Asp	cca Pro	gtc Val	aac Asn	ttg Leu 25	cag Gln	aaa Lys	ttc Phe	tct Ser	tac Tyr 30	act Thr	agt Ser	gag Glu	gat Asp	gag Glu 35	151
gcc Ala	tgg Trp	aag Lys	acg Thr	tac Tyr 40	cta Leu	gaa Glu	aac Asn	ccg Pro	ttg Leu 45	aca Thr	gct Ala	gcc Ala	aca Thr	aag Lys 50	gcc Ala	199
atg Met	atg Met	aga Arg	gtc Val 55	aat Asn	gga Gly	gat Asp	gat Asp	gac Asp 60	agt Ser	gtt Val	gcg Ala	gcc Ala	ttg Leu 65	agc Ser	ttc Phe	247
ctc Leu	tat Tyr	gat Asp 70	tac Tyr	tac Tyr	atg Met	ggt Gly	ccc Pro 75	aag Lys	gag Glu	aag Lys	cgg Arg	ata Ile 80	ttg Leu	tcc Ser	tcc Ser	295
agc Ser	act Thr 85	eja aaa	gly	agg Arg	aat Asn	gac Asp 90	Glņ Glņ	gga Gly	aag Lys	agg Arg	tac Tyr 95	tac Tyr	cat His	ggc	atg Met	343
gaa Glu 100	tat Tyr	gag Glu	acg Thr	gac Asp	ctc Leu 105	act Thr	ccc Pro	ctt Leu	gaa Glu	agc Ser 110	ccc Pro	aca Thr	cac His	ctc Leu	atg Met 115	391
aaa Lys	ytc Xaa	ctg Leu	aca Thr	gag Glu 120	aac Asn	gtg Val	tct Ser	gga Gly	acc Thr 125	cca Pro	gag Glu	tac Tyr	cca Pro	gat Asp 130	ttg Leu	439
ctc Leu	aag Lys	aag Lys	aat Asn 135	aac Asn	ctg Leu	atg Met	agc Ser	ttg Leu 140	gag Glu	Gly	gcc Ala	ttg Leu	ccc Pro 145	acc Thr	cct Pro	487
ggc	aag Lys	gca Ala 150	gct Ala	ccc Pro	ctc Leu	cct Pro	gca Ala 155	ggc	ccc Pro	agc Ser	aag Lys	ctg Leu 160	gag Glu	gcc Ala	Gly	<b>53</b> 5
tct Ser	gtg Val 165	gac Asp	agc Ser	tac Tyr	ctg Leu	tta Leu 170	ccc Pro	acy Xaa	act Thr	gat Asp	atg Met 175	tat Tyr	gat Asp	aat Asn	Gly	583
tcc Ser 180	ctc Leu	aac Asn	tcc Ser	ttg Leu	ttt Phe 185	gag Glu	agc Ser	att Ile	cat His	999 Gly 190	gtg Val	ccg Pro	ccc Pro	aca Thr	cag Gln 195	631
cgc Arg	tgg Trp	cag Gln	cca Pro	gac Asp 200	agc Ser	acc Thr	ttc Phe	aaa Lys	gat Asp 205	gac Asp	cca Pro	cag Gln	gag Glu	tcg Ser 210	atg Met	679
ctc Leu	ttc Phe	cca Pro	gat Asp 215	atc Ile	ctg Leu	aaa Lys	acc Thr	tcc Ser 220	ccg Pro	gaa Glu	ccc Pro	cca Pro	tgt Cys 225	cca Pro	gag Glu	727
gac	tac	ccc	agc	ctc	aaa	agt	gac	ttt	gaa	tac	acc	ctg	ggc	tcc	ccc	775

Asp	Tyr	Pro 230	Ser	Leu	Lys	Ser	Asp 235	Phe	Glu	Tyr	Thr	Leu 240	Gly	Ser	Pro	
					aag Lys											823
		_			ccc Pro 265	_		_				_				871
					tcc Ser											919
	_			Lys	gtc Val		_		_	_	_			_	cac His	. 967
			Arg		ccc Pro			Lys								1015
gac	tgc Cys 325	Ьys	gaa Glu	aac Asn	ttc Phe	aac Asn 330	Thr	gtg Val	gag Glu	cac His	att Ile 335	Glu	gag Glu	gtg Val	gcc	1063
	Asn				ttt Phe 345	Val					Glu					1111
			_		ı Cys	_	_		_	Phe			_	_	) : GJÀ 1 A3A	1159
				Pro					lle			Tyr		Су	ggc Gly	1207
			r Gli					s Arg					ı Ile		g atc s Ile	1255
		s. As					u Ar					aA c			g aag g Lys	1303
	n Ph	_		_		l Ly			_		r Se		_		c gtc y Val 435	
					u Se					y As:					c ctt r Leu 0	
cg	g cc	a ga	g ac	t ga	c ct	g ga	g ac	g cc	a cc	c gt	g ct	g tt	c at	c cc	c aat	1447

Arg Pro	Glu	Thr 455	Asp	Leu	Glu	Thr	Pro 460	Pro	Val	Leu	Phe	Ile 465	Pro	Asn	
gtg cad Val His	ttc Phe 470	tcc Ser	agc Ser	ctg Leu	cag Gln	cgc Arg 475	tct Ser	gga Gly	ely aaa	gca Ala	gcc Ala 480	ccc Pro	tcg Ser	gca Ala	1495
gga ccc Gly Pro 485	Ser	agc Ser	tcc Ser	aac Asn	agg Arg 490	ctg Leu	cct Pro	ctg Leu	aag Lys	cgt Arg 495	acc Thr	tgc Cys	tcg Ser	ccc Pro	1543
ttc act Phe Thi 500	gag Glu	gag Glu	ttt Phe	gag Glu 505	cct Pro	ctg Leu	ccc Pro	tcc Ser	aag Lys 510	cag Gln	gcc Ala	aag Lys	gaa Glu	ggc Gly 515	1591
gac cti Asp Lei	cag Gln	aga Arg	gtt Val 520	ctg Leu	ctg Leu	tat Tyr	gtg Val	cgg Arg 525	agg Arg	gag Glu	act Thr	gag Glu	gag Glu 530	gtg Val	1639
ttt gad Phe Asj	gcg Ala	ctc Leu 535	atg Met	ttg Leu	aag Lys	acc Thr	cca Pro 540	gac Asp	ctg Leu	aag Lys	Gly 333	ctg Leu 545	agg Arg	aat Asn	1687
gcg ato Ala Ilo	tct Ser 550	gag Glu	aag Lys	tat Tyr	ej aaa	ttc Phe 555	cct Pro	gaa Glu	gag Glu	aac Asn	att Ile 560	tac Tyr	aaa Lys	gtc Val	1735
tac aag Tyr Lys 56	s Lys	tgc Cys	aag Lys	cga Arg	gga Gly 570	atc Ile	tta Leu	gtc Val	aac Asn	atg Met 575	gac Asp	aac Asn	aac Asn	atc Ile	1783
att cas Ile Gli 580	g cat n His	tac Tyr	agc Ser	aac Asn 585	cac His	gtc Val	gcc Ala	ttc Phe	ctg Leu 590	ctg Leu	gac Asp	atg Met	Gly 999	gag Glu 595	1831
ctg gad Leu As	ggc Gly	aaa Lys	att Ile 600	cag Gln	atc Ile	atc Ile	ctt Leu	aag Lys 605	gag Glu	ctg Leu	taa				1870
<210><211><212><212><213>	8 607 PRT huma	n													
<220><221><222><223>	(117	ī(	117)		atio	n 11	7 st	ands	for	Leu	, or	Phe	-		
<220> <221> <222> <223>	(172	ī(	172)		atio	n 17	2 st	ands	for	Thr	•				
<400> Met Tr		Asn	Ser	Ile	Leu	Pro	Ile	Phe	Leu	Phe	Arg	Ser	Val	Arg	

1				5					10					15	
Leu	Leu	Lys	Asn 20	Asp	Pro '	Val .		Leu 25	Gln	Lys	Phe		Tyr 30	Thr	Ser
Glu	Asp	Glu 35	Ala	Trp	Lys		Tyr 40	Leu	Glu	Asn		Leu 45	Thr	Ala	Ala
Thr	Lуs 50	Ala	Met	Met	Arg	Val 55	Asn	Gly	Asp	Asp	Asp 60	Ser	Val	Ala	Ala
Leu 65	Ser	Phe	Leu	Tyr	Asp 70	Tyr	Tyr	Met	Gly	Pro 75	Lys	Glu	ГÀв	Arg	Ile 80
Leu	Ser	Ser	Ser	Thr 85	Gly	Gly	Arg	Asn	Asp 90	Gln	Gly	ГХE	Arg	Tyr 95	Tyr
His	Gly	Met	Glu 100	Tyr	Glu	Thr	qaA	Leu 105	Thr	Pro	Leu	Glu	Ser 110	Pro	Thr
His	Leu	Met 115	_	Xaa	Leu	Thr	Glu 120	Asn	Val	Ser	Gly	Thr 125	Pro	Glu	Tyr
Pro	Asp 130		Leu	Lys	ГÀв	Asn 135	Asn	Leu	Met	Ser	Leu 140	Glu	Gly	Ala	Leu
Pro 145		Pro	Gly	' Lys	Ala 150	Ala	Pro	Leu	Pro	Ala 155		Pro	Ser	гÀв	Leu 160
Glu	Ala	. Gly	Ser	Val 165	_	Ser	Tyr	Leu	Leu 170		Xaa	Thr	Asp	Met 175	Tyr
Asp	Asn	Gly	/ Ser 180		Asn	Ser	Leu	Phe 185		Ser	Ile	His	Gly 190	Val	Pro
Pro	Thr	Glr 1,95	_	Trp	Gln	Pro	200		Thr	Phe	. Lys	205		Pro	Gln
Glu	210		: Lev	ı Phe	Pro	Asp 215		. Lev	ı Lys	Thr	220		Glu	Pro	Pro
Cys 225		Gl:	u Asj	о Туг	230		Lev	ı Lys	s Sei	235		: Glv	туг	Thr	Leu 240
Gly	, Sei	r Pro	o Ly	s Ala 24	_	His	s Ile	≥ Ly:	250 250		/ Glu	ı Sei	r Pro	259	Ala
Ty	r Lei	u As:	n Ly 26		y Glr	) Phe	∋ Туг	26		l Th	r Lei	ı Arç	270		o Ala
Gl	y Gl	y Ly 27		у Le	u Ala	a Let	u Se: 28		r Ası	n Ly	в Va:	l Ly: 28:		c Vai	l Val
Me	t Va		l Ph	e As	p Ası	1 Gl		s Va	l Pr	o Va	1 Gl		n Lei	ı Ar	g Phe

Trp Lys His Trp His Ser Arg Gln Pro Thr Ala Lys Gln Arg Val Ile Asp Val Ala Asp Cys Lys Glu Asn Phe Asn Thr Val Glu His Ile Glu 330 Glu Val Ala Tyr Asn Ala Leu Ser Phe Val Trp Asn Val Asn Glu Glu Ala Lys Val Phe Ile Gly Val Asn Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Val Lys Gly Val Pro Leu Asn Leu Gln Ile Asp Thr Tyr 375 Asp Cys Gly Leu Gly Thr Glu Arg Leu Val His Arg Ala Val Cys Gln 395 Ile Lys Ile Phe Cys Asp Lys Gly Ala Glu Arg Lys Met Arg Asp Asp 410 Glu Arg Lys Gln Phe Arg Arg Lys Val Lys Cys Pro Asp Ser Ser Asn Ser Gly Val Lys Gly Cys Leu Leu Ser Gly Phe Arg Gly Asn Glu Thr 445 Thr Tyr Leu Arg Pro Glu Thr Asp Leu Glu Thr Pro Pro Val Leu Phe 455 Ile Pro Asn Val His Phe Ser Ser Leu Gln Arg Ser Gly Gly Ala Ala 470 Pro Ser Ala Gly Pro Ser Ser Ser Asn Arg Leu Pro Leu Lys Arg Thr 490 Cys Ser Pro Phe Thr Glu Glu Phe Glu Pro Leu Pro Ser Lys Gln Ala 505 Lys Glu Gly Asp Leu Gln Arg Val Leu Leu Tyr Val Arg Arg Glu Thr 520 Glu Glu Val Phe Asp Ala Leu Met Leu Lys Thr Pro Asp Leu Lys Gly Leu Arg Asn Ala Ile Ser Glu Lys Tyr Gly Phe Pro Glu Glu Asn Ile Tyr Lys Val Tyr Lys Lys Cys Lys Arg Gly Ile Leu Val Asn Met Asp Asn Asn Ile Ile Gln His Tyr Ser Asn His Val Ala Phe Leu Leu Asp Met Gly Glu Leu Asp Gly Lys Ile Gln Ile Ile Leu Lys Glu Leu

```
<210>
       9
<211>
       3113
<212>
       DNA
<213>
       murine
<220>
<221>
       misc feature
<222>
       (2634)..(2634)
<223> n = any nucleotide
<220>
<221> misc_feature
       (2968) . . (2968)
<222>
       n = any nucleotide
```

<400> 9 gttectecat gggtteettg agtteetgac atggetteec ttgatgatga actgtgtgae 60 ctaaacagca taccaaatgt gacggagcag cccctcattt ctgctggaga aaacagggta 120 caagtgctga aaaacgtgcc cttcaacatc gtcctccccc atagcaacca gctgggcatt 180 gataagagag gccatctgac agctcccgat acaacagtca ctgtctccat agcgaccatg 24Ò cctacccact ccatcaagac agaaatccag ccgcacggct ttgctgtggg aatccctcca 300 360 geogtgtace actetgagee cacegaaege gtggtggttt ttgaceggag ceteageaet gatcagttca gctctggcac tcagcccccc aatgctcagc ggaggactcc agactccacc 420 tteteegaga eetteaagga gggegtteag gaggttttet teecetegga acteageett 480 eggatgeegg geatgaatte agaggaetat gtetttgaea atgtttetgg gaacaaettt 540 600 gagtataccc tggaagcctc caagtcactg cggcagaagc aaggggacag cactatgaca tacctgaata aaggccagtt ctatcctgtc accttaaagg aaggaagcag caatgaaggg .660 atteaceace etateageaa agttegaagt gtgateatgg tggtttttgc tgaagacaaa 720 agcagagaag accagetgag acaetggaag taetggeaet ceegteagea caeggecaaa 780 cagaggtgca ttgacattgc tgactacaaa gaaagtttca acactatcag caacattgag 840 gagatagett ataacgecat tteetteacg tgggacatea atgatgagge aaaggtette 900 atototgtga actgottgag cacagattto tottotoaga agggtgtgaa gggottgoca 960 ctcaacattc aaatcgacac atacagctat aacaaccgca gcaacaagcc ggttcaccgg 1020 gcctactgcc agataaaggt cttctgcgac aagggagctg aaaggaaaat tcgggatgaa 1080 gaacgaaaac agagcaagag aaaagtgtct gacgttaaag tgcagctgct tccctcacac 1140 aaacggacag acatcacagt gttcaagccc ttcctggacc tcgacactca gcctgtcctc 1200 ttcattccgg acgtgcattt taccaacctg cagcggggca gtcatgttct ttccctcccc 1260 tctgaagaac tggaaggtga aggctctgtc ttgaaaagag ggccattcgg aaccgaagat 1320 gactttggag ttcctcctcc tgctaagctg actcggacag aagaacccaa gagagtgctg 1380 ctctatgtcc gaaaggaatc agaagaagtc ttcgacgccc tgatgctcaa gacgccgtct 1440 ttgaagggcc tgatggaggc aatttcagac aagtatgatg tcccccatga caagattggg 1500 aaaatattta agaagtgcaa aaaagggatc ctcgtgaaca tggacgacaa cattgtgaag 1560 cactactcca atgaggacac cttccagctg cagatagagg aagccggcgg ctcgtacaag 1620 ctcaccctga cagagattta aaggggcagg ggtggggggc gctcggctcc caggcgtggg 1680 aattcagtga aagtgttcca gctgagaagc ccaggcacct accctgcaga accttaaata 1740 tcagggaagg aacctttcac gtaggaaatg gcgctgtgta taccgtgctg tgttgatgtt 1800 ttettttgga tagaaateea tgtgttgttt tgttgttgtt gtttgaattt etgatgtget 1860 tagaaagcga agcatgagaa ctttgtaccg gatctaagag accatgggac cgtttgggtt 1920 acctgctcca ctacctgtca aagtctgcct gtgtccataa gagtggtggg ctactggctg 1980 gcgagagagg ggaaggcagt agcttgtctt tgaggctttt gtgttctcgc ctgacctcag 2040 tctaactctg actgccttga ggagtgggcc cagccctcag caataaaggg ctaagccttc 2100 tecetecace tetectecag tgtttaetaa atagggtgea tteetggaae etttteeege 2160 aacttccctt ggacatgtgg actgcctttc tgatgaagaa cttgcgtgag tgacagtgtg 2220 aagttagete tgttaaaget gegttgtata taagtgeaat atetttttga aggtetgeet 2280 gtaaatgtgt acatatatgt ctgatataaa tatataatat ataaatgcgg tgtctgtgta 2340 cagatagtga aggcgagcag gaagatctac cttgaaatcc ctcttagaga agaggttaag 2400 ttattattga taatgtggac caagcaggta gaacgctgtt ttcccaaaaa caagcaagtg 2460 ttccctagca tagcaaaaag ccatctcatg tggcagagcc atctgctctt gcgaatgttg 2520 teacegtgtg ggtttetgca ecetgagtgg agetaatgga agaetggaet geagetaeta 2580 tatgaggtgt gtgtgcaggt gtcagccaag ctgtgcccat gcagagactc agcngtgtca 2640 tgagccagcg attcaaacca aaatgggccg attctacaag gccatgtttc agagcttcca 2700 agcatcaget accetetett teaacteeaa eecattcate aatttacata acteteecae 2760 ' gggaatgttt tgtgcacact taaatattta agaacaaaac gaaactttac aatgtaaytt 2820

tata	atgaa	at co	etgta	aacag	aaa	ataca	aatt	gcg	ggtti	cct	ttagg	gttc	ag g	gaact	cagaa
tagg	tcati	tt gi	tatg	agtag	g ga	ttgti	tagc	ggt	atac	gta	rgtta	aaaa	ag ta	actci	taatg
aagt	atgt	ga a	caaa	atago	e tg	gttt	tnta	aga	tacg	gga	tacg	ggte	at a	taac	aatat
tttc	tatt	tt g	tttt	atgaa	a at	cage	ttta	ctt	gttt	taa	ttgt	atca	tt g	aaca	tgtgt
ttta	aacc	aa a	ggga	ttgaa	a tt	ttat	atgt	cta	tttc	aaa	aaaa	aaaa	aa a	aa	
<210 <211 <212 <213	> .5 > P		e					•		-					
<400 Met 1		0 Ser		Asp : 5	Asp	Glu	Leu	Cys	Asp 10	Leu	Asn	Ser	Ile	Pro 15	Asn
Val	Thr	Glu	<b>Gln</b> 20	Pro	Leu	Ile	Ser	Ala 25	Gly	Glu	Asn	Arg	Val 30	Gln	Val .
Leu	ГÀв	Asn 35	Val	Pro	Phe	Asn	Ile 40	Val	Leu	Pro	His	Ser 45	Asn	Gln	Leu
Gly	Ile 50	Авр	Lys	Arg	Gly	His 55	Leu	Thr	Ala	Pro	Asp 60	Thr	Thr	Val	Thr
Val 65	Ser	Ile	Ala	Thr	Met 70	Pro	Thir	His	Ser	Ile 75	Lys	Thr	Glu	Ile	Gln . 80
Pro	His	Gly	Phe	Ala 85	Val	Gly	Ile	Pro	Pro 90	Ala	Val	Týr	His	Ser 95	Glu
Pro	Thr	Glu	Arg 100	Val	Val	Val	Phe	Asp 105	Arg	Ser	Leu	Ser	Thr 110	Asp	Gln
Phe	Ser	Ser 115	Gly	Thr	Gln	Pro	Pro 120	Asn	Ala	Gln	Arg	Arg 125	Thr	Pro	Asp
Ser	Thr 130	Phe	Ser	Glu	Thr	Phe 135	Lys	Glu	Gly	Val	Gln 140	Glu	Val	Phe	Phe
Pro 145	Ser	Glu	Leu	Ser	Leu 150	Arg	Met	Pro	Gly	Met 155		Ser	Glu	Asp	Tyr 160
Val	Phe	Asp	Asn	Val 165	Ser	Gly	Asn	Asn	Phe 170	Glu	Tyr	Thr	Leu	Glu 175	Ala
Ser	Lys	Ser	Leu 180	Arg	Gln	Lys	Gln	Gly 185		Ser	Thr	Met	Thr 190		Leu
Asn	Lys	Gly 195		Phe	Tyr	Pro	Val 200		Leu	Lys	Glu	Gly 205		Ser	Asn

Glu Gly Ile His His Pro Ile Ser Lys Val Arg Ser Val Ile Met Val 215 Val Phe Ala Glu Asp Lys Ser Arg Glu Asp Gln Leu Arg His Trp Lys Tyr Trp His Ser Arg Gln His Thr Ala Lys Gln Arg Cys Ile Asp Ile 250 Ala Asp Tyr Lys Glu Ser Phe Asn Thr Ile Ser Asn Ile Glu Glu Ile 260 Ala Tyr Asn Ala Ile Ser Phe Thr Trp Asp Ile Asn Asp Glu Ala Lys 280 Val Phe Ile Ser Val Asn Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Val Lys Gly Leu Pro Leu Asn Ile Gln Ile Asp Thr Tyr Ser Tyr 315 Asn Asn Arg Ser Asn Lys Pro Val His Arg Ala Tyr Cys Gln Ile Lys Val Phe Cys Asp Lys Gly Ala Glu Arg Lys Ile Arg Asp Glu Glu Arg Lys Gln Ser Lys Arg Lys Val Ser Asp Val Lys Val Gln Leu Leu Pro Ser His Lys Arg Thr Asp Ile Thr Val Phe Lys Pro Phe Leu Asp Leu 375 Asp Thr Gln Pro Val Leu Phe Ile Pro Asp Val His Phe Thr Asn Leu Gln Arg Gly Ser His Val Leu Ser Leu Pro Ser Glu Glu Leu Glu Gly Glu Gly Ser Val Leu Lys Arg Gly Pro Phe Gly Thr Glu Asp Asp Phe 425 Gly Val Pro Pro Pro Ala Lys Leu Thr Arg Thr Glu Glu Pro Lys Arg Val Leu Leu Tyr Val Arg Lys Glu Ser Glu Glu Val Phe Asp Ala Leu Met Leu Lys Thr Pro Ser Leu Lys Gly Leu Met Glu Ala Ile Ser Asp 470 475 Lys Tyr Asp Val Pro His Asp Lys Ile Gly Lys Ile Phe Lys Lys Cys Lys Lys Gly Ile Leu Val Asn Met Asp Asp Asn Ile Val Lys His Tyr 505

Ser Asn Glu Asp Thr Phe Gln Leu Gln Ile Glu Glu Ala Gly Gly Ser 515 520 Tyr Lys Leu Thr Leu Thr Glu Ile 530 <210> 11 <211> 3452 <212> DNA <213> murine <220> <221> misc\_feature (2973) . . (2973) <222> <223> n = any nucleotide <220> <221> misc\_feature <222> (3307)..(3307) <223> n = any nucleotide cgccgctccg gacccaccgc ctgccgccgc gcgccgcccg ccgccgcctc ctccccccgg 60 ategggtgta ctgtcccaac ccgaaagtcc agttctgcgg cccggcagcg gcgagcgagc 120 180 gcgatgacac aggagtacga caacaaaagg cccgtgctgg tacttcagaa tgaagccctc 240 tacccacage ggcgctccta taccagtgag gatgaageet ggaagtegtt cetggaaaac 300 ceteteactg eggeaaceaa agegatgatg ageateaacg gagacgaaga cagegegget gegetgggee tgetetatga etaetaeaag gteeceagag agegeeggte ateageegta 360 420 aagcccgagg gagagcaccc agagccagag cacagcaaaa gaaacagcat accaaatgtg acggagcagc ccctcatttc tgctggagaa aacagggtac aagtgctgaa aaacgtgccc 480 ttcaacatcg tcctccccca tagcaaccag ctgggcattg ataagagagg ccatctgaca 540 getecegata caacagteac tgtetecata gegaceatge etaeceacte cateaagaca 600 gaaatccagc cgcacggett tgctgtggga atccctccag ccgtgtacca ctctgagccc 660 720 accgaacgcg tggtggtttt tgaccggagc ctcagcactg atcagttcag ctctggcact 780 cagccccca atgetcageg gaggaeteca gaetecaeet tetcegagae etteaaggag 840 ggcgttcagg aggttttctt cccctcggaa ctcagccttc ggatgccggg catgaattca

gaggactatg tetttgacaa tgtttetggg aacaaetttg agtataeeet ggaageetee

aagtcactgc ggcagaagca aggggacagc actatgacat acctgaataa aggccagttc

tatcctgtca ccttaaagga aggaagcagc aatgaaggga ttcaccaccc tatcagcaaa

900

960

gttcgaagtg tgatcatggt ggtttttgct gaagacaaaa gcagagaaga ccagctgaga 1080 1140 cactggaagt actggcactc ccgtcagcac acggccaaac agaggtgcat tgacattgct gactacaaag aaagtttcaa cactatcagc aacattgagg agatagctta taacgccatt 1200 1260 teetteaegt gggacateaa tgatgaggca aaggtettea tetetgtgaa etgettgage 1320 acagatttet etteteagaa gggtgtgaag ggettgeeac teaacattea aategacaca tacagotata acaacegeag caacaageeg gttcaceggg cetactgeca gataaaggte 1380 ttctgcgaca agggagctga aaggaaaatt cgggatgaag aacgaaaaca gagcaagaga 1440 aaagtgtctg acgttaaagt gcagctgctt ccctcacaca aacggacaga catcacagtg 1500 ttcaagecet teetggacet egacacteag cetgteetet teatteegga egtgeatttt 1560 accaacctgc agcggggcag tcatgttctt tccctccct ctgaagaact ggaaggtgaa 1620 ggctctgtct tgaaaagagg gccattcgga accgaagatg actttggagt tcctcctcct 1680 gctaagctga ctcggacaga agaacccaag agagtgctgc tctatgtccg aaaggaatca 1740 gaagaagtet tegaegeest gatgeteaag aegeegtett tgaagggeet gatggaggea 1800 1860 atttcagaca agtatgatgt cccccatgac aagattggga aaatatttaa gaagtgcaaa aaagggatcc tcgtgaacat ggacgacaac attgtgaagc actactccaa tgaggacacc 1920 1980 ttccagctgc agatagagga agccggcggc tcgtacaagc tcaccctgac agagatttaa 2040 aggggcaggg gtggggggcg ctcggctccc aggcgtggga attcagtgaa agtgttccag 2100 ctgagaagcc caggcaccta ccctgcagaa ccttaaatat cagggaagga acctttcacg taggaaatgg cgctgtgtat accgtgctgt gttgatgttt tcttttggat agaaatccat 2160 gtgttgtttt gttgttgttg tttgaatttc tgatgtgctt agaaagcgaa gcatgagaac 2220 tttgtaccgg atctaagaga ccatgggacc gtttgggtta cctgctccac tacctgtcaa 2280 agtetgeetg tgteeataag agtggtggge taetggetgg cgagagaggg gaaggeagta 2340 gettgtettt gaggettttg tgttetegee tgaceteagt etaactetga etgeettgag 2400 gagtgggccc agccctcagc aataaagggc taagccttct ccctccacct ctcctccagt 2460 gtttactaaa tagggtgcat tcctggaacc ttttcccgca acttcccttg gacatgtgga 2520 etgeetttet gatgaagaae ttgegtgagt gaeagtgtga agttagetet gttaaagetg 2580 cgttgtatat aagtgcaata tctttttgaa ggtctgcctg taaatgtgta catatatgtc 2640 tgatataaat atataatata taaatgoggt gtotgtgtac agatagtgaa ggogagcagg 2700

aagatctacc	ttgaaatccc	tcttagagaa	gaggttaagt	tattattgat	aatgtggacc	2760
aagcaggtag	aacgctgttt	tcccaaaaac	aagcaagtgt	tccctagcat	agcaaaaagc	2820
catctcatgt	ggcagagcca	tetgetettg	cgaatgttgt	.caccgtgtgg	gtttctgcac	2880
cctgagtgga	gctaatggaa	gactggactg	cagctactat	atgaggtgtg	tgtgcaggtg	2940
tcagccaagc	tgtgcccatg	cagagactca	gengtgteat	gagccagcga	ttcaaaccaa	3000
aatgggccga	ttctacaagg	ccatgtttca	gagettecaa	gcatcagcta	ccgtgtgttt	3060
gaactggaag	gcattcatga	atttacataa	ctgtggcagg	ggaatgtttt	gtgcacactt	3120
aaatatttaa	gaacaaaacg	aaactttaca	atgtaayttt	ataatgaatc	ctgtaacaga	3180
aatacaattg	cgggtttctt	taggttcagg	gaactagaat	aggtcatttg	tatgagtagg	3240
attgttagcg	gtatacgtar	gttaaaaagt	actctaatga	agtatgtgaa	caaaatagct	3300
ggttttntaa	gatacgggat	acgggtcata	taacaatatt	ttctattttg	ttttatgaaa	3360
tcagctttac	ttgttttaat	tgtatcattg	aacatgtgtt	ttaaaccaaa	gggattgaat	3420
tttatatgtc	tatttcaaaa	aaaaaaaaa	aa			3452
<210> 12						

<211> 618

<212> PRT

<213> murine

<400> 12

Met Thr Gln Glu Tyr Asp Asn Lys Arg Pro Val Leu Val Leu Gln Asn 1 5 10 15.

Glu Ala Leu Tyr Pro Gln Arg Arg Ser Tyr Thr Ser Glu Asp Glu Ala 20 25 30

Trp Lys Ser Phe Leu Glu Asn Pro Leu Thr Ala Ala Thr Lys Ala Met 35 40 45

Met Ser Ile Asn Gly Asp Glu Asp Ser Ala Ala Leu Gly Leu Leu 50 55 60

Tyr Asp Tyr Tyr Lys Val Pro Arg Glu Arg Arg Ser Ser Ala Val Lys 65 70 75 80

Pro Glu Gly Glu His Pro Glu Pro Glu His Ser Lys Arg Asn Ser Ile 85 90 95

Pro Asn Val Thr Glu Gln Pro Leu Ile Ser Ala Gly Glu Asn Arg Val

Gln Val Leu Lys Asn Val Pro Phe Asn Ile Val Leu Pro His Ser Asn 115 120 125

Gln Leu Gly Ile Asp Lys Arg Gly His Leu Thr Ala Pro Asp Thr Thr 130 135 140

Val Thr Val Ser Ile Ala Thr Met Pro Thr His Ser Ile Lys Thr Glu
145 150 150 160

Ile Gln Pro His Gly Phe Ala Val Gly Ile Pro Pro Ala Val Tyr His 165 170 175

Ser Glu Pro Thr Glu Arg Val Val Val Phe Asp Arg Ser Leu Ser Thr 180 185 190

Asp Gln Phe Ser Ser Gly Thr Gln Pro Pro Asn Ala Gln Arg Arg Thr 195 200 205

Pro Asp Ser Thr Phe Ser Glu Thr Phe Lys Glu Gly Val Gln Glu Val 210 215 220

Phe Phe Pro Ser Glu Leu Ser Leu Arg Met Pro Gly Met Asn Ser Glu 225 230 235 240

Asp Tyr Val Phe Asp Asn Val Ser Gly Asn Asn Phe Glu Tyr Thr Leu 255 255

Glu Ala Ser Lys Ser Leu Arg Gln Lys Gln Gly Asp Ser Thr Met Thr 260 265 270

Tyr Leu Asn Lys Gly Gln Phe Tyr Pro Val Thr Leu Lys Glu Gly Ser 275 280 285

Ser Asn Glu Gly Ile His His Pro Ile Ser Lys Val Arg Ser Val Ile 290 295 300

Met Val Val Phe Ala Glu Asp Lys Ser Arg Glu Asp Gln Leu Arg His 305 310 315 320

Trp Lys Tyr Trp His Ser Arg Gln His Thr Ala Lys Gln Arg Cys Ile 325 330 335

Asp Ile Ala Asp Tyr Lys Glu Ser Phe Asn Thr Ile Ser Asn Ile Glu 340 345

Glu Ile Ala Tyr Asn Ala Ile Ser Phe Thr Trp Asp Ile Asn Asp Glu 355 360 365

Ala Lys Val Phe Ile Ser Val Asn Cys Leu Ser Thr Asp Phe Ser Ser 370 375 380

Gln Lys Gly Val Lys Gly Leu Pro Leu Asn Ile Gln Ile Asp Thr Tyr 385 390 395 400

Ser Tyr Asn Asn Arg Ser Asn Lys Pro Val His Arg Ala Tyr Cys Gln
405
415

Ile Lys Val Phe Cys Asp Lys Gly Ala Glu Arg Lys Ile Arg Asp Glu

			420					425					430			
Glu	Arg	Lув 435	Gln	Ser	Lys	Arg	Lys 440	Val	Ser	Asp	Val	Lys 445	Val	Gln	Leu	
Leu	Pro 450	Ser	His	ГÀв	Arg	Thr 455	qaA	Ile	Thr	Val	Phe 460	Lys	Pro	Phe	Leu	٠
Asp 465		Asp	Thr	Gln	Pro 470	Val	Leu	Phe	Ilė	Pro 475	Asp	Val	His	Phe	Thr 480	
Asn	Leu	Gln	Arg	Gly 485		His	Val	Leu	Ser 490	Leu	Pro	Ser	Glu	Glu 495	Leu	
Glu	Gly	Glu	Gly 500		Val	Leu	Lys	Arg 505	Gly	Pro	Phe	Gly	Thr 510	Glu	Asp.	
Asp	Phe	Gly 515		Pro	Pro	Pro	Ala 520		Leu	Thr	Arg	525	Glu	Glu	Pro	
Lys	530		Leu	Let	Tyr	Val 535	Arg	j Lys	s Glu	ser	540	ı Glu	val	. Phe	e Asp	
Ala 549		ı Met	. Lev	ı Lys	550		Se:	r, Leı	і Гу	555	y Let	u Met	: Glu	a Ala	11e 560	
Se	c As	, Ly	в Туг	r Ası 56!		l Pro	Hi:	s Asj	р <b>L</b> ys 57	g Ile	e Gl	Ty	s Ile	9 Phe 57!	e Lys 5	
Ly	в Су	s Ly	s Ly:		y Ile	e Lei	u Va	1 As: 58	n Me	t As	p As	p As	n Ile 59	e Vai	l Lys	
нi	в Ту	r Se 59		n Gl	u As	p Th	r Ph 60	e Gl O	n Le	u Gl	n Il	e G1 60	u Gl <sup>.</sup> 5	u Al	a Gly	
G1	y Se 61		r Ly	s Le	u Th	r Le 61		ır Gl	u Il	.e	•					
<2 <2	210> 211> 212> 213>	13 219 DNA mus	4			٠										
<4 C9	300>	13 gggca	a ggt	caga	actt	gaaa	agtc	cag 1	ttte	accag	ga g	gctga	aggct	cca	aggaaaag	g 60
g	ggag	cgagʻ	t to	attg	gatc	aaad	catg	tca (	caag	agtc	gg a	caata	aataa	a aag	gactagte	g 120
g	cctt	agtg	c cc	atgc	ccag	tga	ccct	ccc	ttca	acac	cc g	aaga	gcct	a ca	caagtga	g 18
g	atga	ggcc	t gg	aagt	cata	tot	ggag	aac	cccc	tgac	tg c	ggcc	acca	a gg	cgatgat	g 24
a	gcat	caac	g gg	gacg	agga	cag	tgct	gcc	gccc	tggg	cc t	gctc	tatg	a ct	actacaa	g 30
q	ttcc	tcga	g ac	aaga	gact	tct	gtct	gtg	agca	aagc	aa g	tgac	agcc	a ag	aagacca	.g 36

gataaaagaa actgeettgg caccagtgaa geecagatea atttgagegg aggegagaae 420 agagtgcagg ttctgaagac tgtcccggtg aacctctgtc taagtcaaga ccacatggag 480 aattegaage gegageagta cagtgtatee ateaecgaga getetgeegt cateecegtg 540 teaggeatea eegtggtgaa ageegaggat tteacacegg tgtteatgge geeceeggtg 600 cactatecee gegeggacag tgaggageag egegtggtta tetttgaaca gaeteagtae 660 gacctgccct ccatagccag ccacagctcc tatctcaagg acgaccagcg cagcacgccg 720 780 gacagcacct acagcgagag ctttaaggac ggcgcctcgg agaaatttcg gagtacttct gttggtgctg acgagtatac atatgaccag acgggaagtg gtacatttca gtacaccctg 840 900 gaagccacca aatctctccg tcagaaacag ggggagggcc ccatgaccta cctcaacaaa ggacaattet atgecataac acteagtgag actggagaca acaaatgett cegacaecee 960 atcagcaaag tcaggagtgt ggtgatggtg gtctttagtg aagacaaaaa ccgagatgag 1020 cagetgaaat aetggaagta etggeaetee eggeageaea etgeeaagea gagggteett 1080 gacattgctg attacaagga gagcttcaac accatcggga acattgaaga gatcgcatac 1140 1200 aatgetgttt cetteacetg ggatgtgaae gaggaggeaa agatttttat caeegtgaat tgcctgagta cagatttctc ctcccaaaag ggtgtaaaag gacttcccct gatgattcag 1260 atcgacaegt acagctacaa caacegcage aataaaccea tecacagage atactgecag 1320 1380 atcaaggtct tctgtgacaa gggagcagaa agaaaaatcc gggatgaaga gagaaagcag aacaggaaga aagggaaggg ccaggcctct caagcccagt gcaacaactc ctctgatggg 1440 aagatggccg ccataccgtt acagaagaag agtgacatca cgtacttcaa aaccatgccc 1500 gacetgeact cacageetgt getetteata ceagatgtte actttgeaaa eetacagagg 1560 accggacagg tttattacaa cacagacgat gagcgagaag gcagcagcgt ccttgttaag 1620 eggatgttea ggeceatgga agaggagttt ggtecaacae egtetaagea gateaaagaa 1680 gaaaacgtaa aacgagtgct tttatatgtg aggaaggaga acgatgacgt cttcgatgct 1740 ctgatgctga aatcacccac ggtgaagggt ctgatggaag cgctgtctga gaagtatggg 1800 ctgccagtgg agaaaatcac aaagctttat aagaagagca aaaagggcat cctggtcaac 1860 atggatgaca acatcattga gcactattca aatgaggaca ccttcatcct caacatggag 1920 agcatggtgg aaggetteaa gateacgetg atggagatet gageeetggg tgteeeeteg 1980 ataggagett ttggtataet eetteetggg agagatggga tetetgeege eecaggaeet 2040

ggagacccac ccatctcact cacctctcaa gactgttaca agactgctgg gaagggggc	2100
agggcccaag gcccagtaat ggacttcctt caactcttcc acttgctccc tatggagctg	2160
aageetgage eceteageaa atttettete gtgee	2195
<210> 14 <211> 625 <212> PRT <213> murine	
<pre>&lt;400&gt; 14 Met Ser Gln Glu Ser Asp Asn Asn Lys Arg Leu Val Ala Leu Val Pro 15 1</pre>	•
Met Pro Ser Asp Pro Pro Phe Asn Thr Arg Arg Ala Tyr Thr Ser Glu 20 25 30	
Asp Glu Ala Trp Lys Ser Tyr Leu Glu Asn Pro Leu Thr Ala Ala Thr 45	
Lys Ala Met Met Ser Ile Asn Gly Asp Glu Asp Ser Ala Ala Ala Leu 50 55	
Gly Leu Leu Tyr Asp Tyr Tyr Lys Val Pro Arg Asp Lys Arg Leu Leu 65 70 . 75 80	
Ser Val Ser Lys Ala Ser Asp Ser Gln Glu Asp Gln Asp Lys Arg Asn 95 90	
Cys Leu Gly Thr Ser Glu Ala Gln Ile Asn Leu Ser Gly Gly Glu Asn 100 105 110	
Arg Val Gln Val Leu Lys Thr Val Pro Val Asn Leu Cys Leu Ser Gln 115 120 125	
Asp His Met Glu Asn Ser Lys Arg Glu Gln Tyr Ser Val Ser Ile Thr 130 135 140	
Glu Ser Ser Ala Val Ile Pro Val Ser Gly Ile Thr Val Val Lys Ala 145 150 155 160	
Glu Asp Phe Thr Pro Val Phe Met Ala Pro Pro Val His Tyr Pro Arg 165 170 175	
Ala Asp Ser Glu Glu Gln Arg Val Val Ile Phe Glu Gln Thr Gln Tyr 180 185 190	•
Asp Leu Pro Ser Ile Ala Ser His Ser Ser Tyr Leu Lys Asp Asp Gln 195 200 205	•
Arg Ser Thr Pro Asp Ser Thr Tyr Ser Glu Ser Phe Lys Asp Gly Ala 210 215 220	
Ser Glu Lys Phe Arg Ser Thr Ser Val Gly Ala Asp Glu Tyr Thr Tyr 225 230 235	: )

Asp Gln Thr Gly Ser Gly Thr Phe Gln Tyr Thr Leu Glu Ala Thr Lys 250 Ser Leu Arg Gln Lys Gln Gly Glu Gly Pro Met Thr Tyr Leu Asn Lys Gly Gln Phe Tyr Ala Ile Thr Leu Ser Glu Thr Gly Asp Asn Lys Cys 275 Phe Arg His Pro Ile Ser Lys Val Arg Ser Val Val Met Val Val Phe Ser Glu Asp Lys Asn Arg Asp Glu Gln Leu Lys Tyr Trp Lys Tyr Trp 310 His Ser Arg Gln His Thr Ala Lys Gln Arg Val Leu Asp Ile Ala Asp 325 330 Tyr Lys Glu Ser Phe Asn Thr Ile Gly Asn Ile Glu Glu Ile Ala Tyr Asn Ala Val Ser Phe Thr Trp Asp Val Asn Glu Glu Ala Lys Ile Phe 360 Ile Thr Val Asn Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Val Lys Gly Leu Pro Leu Met Ile Gln Ile Asp Thr Tyr Ser Tyr Asn Asn Arg Ser Asn Lys Pro Ile His Arg Ala Tyr Cys Gln Ile Lys Val Phe 405 Cys Asp Lys Gly Ala Glu Arg Lys Ile Arg Asp Glu Glu Arg Lys Gln Asn Arg Lys Lys Gly Lys Gly Gln Ala Ser Gln Ala Gln Cys Asn Asn Ser Ser Asp Gly Lys Met Ala Ala Ile Pro Leu Gln Lys Lys Ser Asp Ile Thr Tyr Phe Lys Thr Met Pro Asp Leu His Ser Gln Pro Val Leu 465 470 Phe Ile Pro Asp Val His Phe Ala Asn Leu Gln Arg Thr Gly Gln Val Tyr Tyr Asn Thr Asp Asp Glu Arg Glu Gly Ser Ser Val Leu Val Lys 505 Arg Met Phe Arg Pro Met Glu Glu Glu Phe Gly Pro Thr Pro Ser Lys Gln Ile Lys Glu Glu Asn Val Lys Arg Val Leu Leu Tyr Val Arg Lys

	530					535					540					
Glu 545	Asn	Asp	Asp	Val	Phe 550	Asp	Ala	Leu	Met	Leu 555	Lys	Ser	Pro	Thr	Val 560	
ГÀЗ	Gly	Leu	Met	Glu 565	Ala	Leu	Ser	Glu	Lys 570	Tyr	Gly	Leu	Pro	Val 575	Glu	
Lys	Ile	Thr	Lys 580	Leu	Tyr	ГÀа	Lys	Ser 585	ГÀв	Lys	Gly	Ile	Leu 590	Val	Asn	
Met	Asp	Asp 595	Asn	Ile	Ile	Glu	His 600		Ser	Asn	Glu	Asp 605	Thr	Phe	Ile	
Leu	Asn 610	Met	Glu	Ser	Met	Val 615		Gly	Phe	Lув	11e 620	Thr	Leu	Met	Glu	
Ile 625																
<21 <21	.1> .2>	15 2831 DNA muri														`,
<22 <22	20> 21> 22> 23>	CDS (200	))	(2008	3)											
<2: <2:	20> 21> 22> 23>	(280	6).	. (28		ide										
<4 ac	00> ctgt	15 gctt	cca	gcca	atc :	agcg	ccac	cg c	agcc	gg <b>g</b> g	a cc	gctg	tcag	caa	aatctca	. 60
ac	atcc	agag	cgc	aacg	tag	agca	aacg	ct t	cccc	gggc	a gg	aagg	gaat	gtc	tgtgtca	120
															gatcgct	
<b>g</b> 9	aacc	tcgg	aga	agga	ag a M 1	et S	cg a er A	at g sn G	aa c lu I	eu A	at t sp P	tc a he A	gg t rg S	er V	tg cgg al Arg .0	232
tt Le	g ct eu Le	g aa u Ly	g aa rs As 15	n As	c cc p Pr	t gt	g ag il Se	gc tt er Pl 20	ie Gl	ig aa in Ly	ıg tt /s Pl	t co le Pr	c ta o Ty 25	r Se	gt aat er Asn	28
G:	ag ga Lu As	ac ga ap G]	u Al	ec to la Ti	g aa	ng ac rs Th	a ta ir Ty 3!	yr Le	eg ga	ag aa lu Aa	ac co an Pi	ct tt co Le 40	eu Th	g go ar Al	et gcc la Ala	32
a T	cc as	aa go ys Al	cc at	et Me	tg ag	ga gt	tc aa	ac g	39 9 ly A	ac ga sp G	ag ga lu G	ag ag	gt gt er Va	g go	ct gct la Ala	37

50 45 ctg agc ttc ctc tac gac tac tat atg ggt ccc aag gag aag cgg ata 424 Leu Ser Phe Leu Tyr Asp Tyr Tyr Met Gly Pro Lys Glu Lys Arg Ile 65 ctg tcc tcc agc act ggt ggc cgg aat gac caa gga aag aag ttc tac 472 Leu Ser Ser Ser Thr Gly Gly Arg Asn Asp Gln Gly Lys Lys Phe Tyr 80 cac age atg gae tat gag deg gat ett ged dec etc gag age ecc aca 520 His Ser Met Asp Tyr Glu Pro Asp Leu Ala Pro Leu Glu Ser Pro Thr 105 cac etc atg aaa ttt ttg aca gag aac gtg tet gga agt eca gae tac 568 His Leu Met Lys Phe Leu Thr Glu Asn Val Ser Gly Ser Pro Asp Tyr 115 110 aca gac cag ctc aag aaa aac aat ctg cta ggc ttg gag ggg gtt cta 616 Thr Asp Gln Leu Lys Lys Asn Asn Leu Leu Gly Leu Glu Gly Val Leu 130 125 ccc acc ccc ggc aag acc aat acc gtc ccc cca ggt ccg agt aaa ctg 664 Pro Thr Pro Gly Lys Thr Asn Thr Val Pro Pro Gly Pro Ser Lys Leu 140 gaa gcc agc tcc atg gac agc tac ctc ttg ccc gcc agt gac ata tat 712 Glu Ala Ser Ser Met Asp Ser Tyr Leu Leu Pro Ala Ser Asp Ile Tyr 165 gac aat ggc tcc ctc aac tca tta ttt gag agc att cat ggg gtt cca 760 Asp Asn Gly Ser Leu Asn Ser Leu Phe Glu Ser Ile His Gly Val Pro 180 ccc aca cag cgc tgg cag cca gac agc acc ttc aaa gat gac cca cag 808 Pro Thr Gln Arg Trp Gln Pro Asp Ser Thr Phe Lys Asp Asp Pro Gln 195 190 gag tot ctg etc ttc cct gat att ctg aag aca tee ccg gac ccc cca 856 Glu Ser Leu Leu Phe Pro Asp Ile Leu Lys Thr Ser Pro Asp Pro Pro 205 tgc cca gag gat tat cca ggc ctc aag agt gac ttt gaa tac acc ctg 904 Cys Pro Glu Asp Tyr Pro Gly Leu Lys Ser Asp Phe Glu Tyr Thr Leu 225 220 ggc tcc ccc aaa gcc att cac atc aaa gca ggg gag tca ccc atg gcc 952 Gly Ser Pro Lys Ala Ile His Ile Lys Ala Gly Glu Ser Pro Met Ala 240 tac ctc aac aag ggt cag ttc tac ccc gtc acc cta cgc acc cca gca 1000 Tyr Leu Asn Lys Gly Gln Phe Tyr Pro Val Thr Leu Arg Thr Pro Ala gga ggg aaa ggc ctc gct ctg tcc tcc agc aaa gtc aag agc gtg gtg 1048 Gly Gly Lys Gly Leu Ala Leu Ser Ser Lys Val Lys Ser Val Val

270	275	280	
atg gtc gtg ttc gat Met Val Val Phe Asp 285	aat gac aag gtc ccc Asn Asp Lys Val Pro 290	gtg gag cag ctg cg Val Glu Gln Leu Ar 295	gt ttc · 1096 cg Phe
tgg agg cac tgg cat Trp Arg His Trp His 300	tcc cgg cag ccc acc Ser Arg Gln Pro Thr 305	gcc aag cag cgc g Ala Lys Gln Arg V 310	tc atc 1144 al Ile 315
	aag gaa aac ttc aac Lys Glu Asn Phe Asn . 325		tt gaa 1192 le Glu 30
gag gtg gcc tat aad Glu Val Ala Tyr Asi 335	gcg ctg tcc ttt gtg Ala Leu Ser Phe Va 340	g tgg aat gtc aac g L Trp Asn Val Asn G 345	gag gaa 1240 Hu Glu
gcc aag gtg ttt at Ala Lys Val Phe Il 350	c ggt gtc aac tgt ct e Gly Val Asn Cys Le 355	g agc aca gac ttc t u Ser Thr Asp Phe : 360	ccc tcg 1288 Ser Ser
	g ggt gtc ccc ctg aa s Gly Val Pro Leu As 370	c ttg caa att gac n Leu Gln Ile Asp 375	acc tat 1336 Thr Tyr
	c act gag cgc ctg gt y Thr Glu Arg Leu Va 385	a cac cgt gct gtc al His Arg Ala Val 390	tgc cag 1384 Cys Gln 395
atc aag atc ttc to	gt gat aag gga gct g ys Asp Lys Gly Ala G 00	ag agg aag atg cgc lu Arg Lys Met Arg 05	gat gat 1432 Asp Asp 410
gaa cgg aag cag t Glu Arg Lys Gln P 415	tt cga agg aag gtc a he Arg Arg Lys Val L 420	ag tgc cca gac tcc ys Cys Pro Asp Ser 425	agt aac 1480 Ser Asn
aat gca gga atc a Asn Ala Gly Ile I 430	ag ggc tgc ctg ctg t ys Gly Cys Leu Leu S 435	ca ggc ttc agg ggc Ser Gly Phe Arg Gly 440	aat gag 1528 Asn Glu
	egg cca gaa act gac o Arg Pro Glu Thr Asp 1 450	ctg gag acc cag cct Leu Glu Thr Gln Pro 455	gtg ttg 1576 Val Leu
	ctg cat ttt tcc agc Leu His Phe Ser Ser 465	cta cag cgc cca gga Leu Gln Arg Pro Gl 470	a ggg gtt 1624 y Gly Val 475
	gga cac agc agc tct Gly His Ser Ser Ser 480	gac agg ctg cct ct Asp Arg Leu Pro Le 485	g aag cga 1672 u Lys Arg 490
acc tgc tca ccc Thr Cys Ser Pro	ttt gct gag gag ttt Phe Ala Glu Glu Phe	gag cct ctt cct to Glu Pro Leu Pro Se	t aaa caa 1720 r Lys Gln

500 505 495 gcc aag gaa gat gac ctt cag aga gtt ctg ttg tat gtg agg agg gag 1768 Ala Lys Glu Asp Asp Leu Gln Arg Val Leu Leu Tyr Val Arg Arg Glu 510 1816 aca gag gag gtg ttt gac gcg ctc atg ttg aag acc ccg gac ctg aag Thr Glu Glu Val Phe Asp Ala Leu Met Leu Lys Thr Pro Asp Leu Lys 535 525 530 1864 gge etg agg aat geg ate tet gag aag tae gge ete eee gag gag aat Gly Leu Arg Asn Ala Ile Ser Glu Lys Tyr Gly Leu Pro Glu Glu Asn 1912 att tgc aaa gtc tac aag aaa tgc aag cga ggc atc ctg gtt aac atg Ile Cys Lys Val Tyr Lys Lys Cys Lys Arg Gly Ile Leu Val Asn Met 565 560 qac aac aac atc atc caa cac tac agc aac cac gtg gcc ttc ctg ctg 1960 Asp Asn Asn Ile Ile Gln His Tyr Ser Asn His Val Ala Phe Leu Leu 580 gac atg ggt gag ctg gac ggc aag atc cag atc atc ctg aag gag cta 2008 Asp Met Gly Glu Leu Asp Gly Lys Ile Gln Ile Ile Leu Lys Glu Leu 590 595 tgagggcccg gcctcaagcg tcccacaccc ggggcccggc tcaagccacg tacaacctct 2068 2128 tetgtgteag etgttaettg aaatgeettt etttgggaaa gaggtetege aageaaceaa ctcggtgatg tccaagccag ggagagacca agaaggttcc aggatctaaa tgtcccaccc 2188 aggetegaac teactecaga getteetgaa ageacecage ceaceggaga gtetgageaa 2248 2308 cacagaccca actgcctgct ttctcttcta agtcccgctg cagaggccct tacaggggac 2368 gggggtcaca ccaccttete tgcagggcta cacccgetgt etcgateggt tetgacgtte 2428 actgtttcct ttctaccaac ttcagaccag agagttctca cactttggcc aaataacttg aaaactcgtg actttcacag cagatgcctt tgtgaggccc ttggagagga aactttctta 2488 ttgacttcct cggcacaaga tgtaagtcac catcatcgag ctgacaggaa caaataccct 2548 2608 tgccacctac tgttgtacac atttcttatt tacagttttc attatgtgat tatatatata tatatgtaag tatatattat gtacatatat gcaacatttt gtatgtccat gttacatttt 2668 tatcatttca aaaatatqta tttcatattt cttgaactat ttttttagct gttattcgat 2728 2788 tatgcatttt gtatatcata gggtttagta ataaaagcct acccatgcac acttaaaaaa aaaaaaaaaa aaatatcnag cttatcgata ccgtcgacct cga 2831

<210> 16 <211> 603 <212> PRT <213> murine

<220>

<221> misc\_feature

<222> (2806)..(2806)

<223> n = any nucleotide

<400> 16

Met Ser Asn Glu Leu Asp Phe Arg Ser Val Arg Leu Leu Lys Asn Asp 1 5 10 15

Pro Val Ser Phe Gln Lys Phe Pro Tyr Ser Asn Glu Asp Glu Ala Trp
20 25 30

Lys Thr Tyr Leu Glu Asn Pro Leu Thr Ala Ala Thr Lys Ala Met Met 35 40 45

Arg Val Asn Gly Asp Glu Glu Ser Val Ala Ala Leu Ser Phe Leu Tyr 50 55 60

Asp Tyr Tyr Met Gly Pro Lys Glu Lys Arg Ile Leu Ser Ser Ser Thr 65 70 75 80

Gly Gly Arg Asn Asp Gln Gly Lys Lys Phe Tyr His Ser Met Asp Tyr 85 90 95

Glu Pro Asp Leu Ala Pro Leu Glu Ser Pro Thr His Leu Met Lys Phe 100 105 110

Leu Thr Glu Asn Val Ser Gly Ser Pro Asp Tyr Thr Asp Gln Leu Lys 115 120 125

Lys Asn Asn Leu Leu Gly Leu Glu Gly Val Leu Pro Thr Pro Gly Lys 130 135 140

Thr Asn Thr Val Pro Pro Gly Pro Ser Lys Leu Glu Ala Ser Ser Met 145 150 155 160

Asp Ser Tyr Leu Leu Pro Ala Ser Asp Ile Tyr Asp Asn Gly Ser Leu 165 170 175

Asn Ser Leu Phe Glu Ser Ile His Gly Val Pro Pro Thr Gln Arg Trp 180 185 190

Gln Pro Asp Ser Thr Phe Lys Asp Asp Pro Gln Glu Ser Leu Leu Phe 195 200 205

Pro Asp Ile Leu Lys Thr Ser Pro Asp Pro Pro Cys Pro Glu Asp Tyr 210 215 220

Pro Gly Leu Lys Ser Asp Phe Glu Tyr Thr Leu Gly Ser Pro Lys Ala 225 230 235 240

Ile His Ile Lys Ala Gly Glu Ser Pro Met Ala Tyr Leu Asn Lys Gly

245 250 255 Gln Phe Tyr Pro Val Thr Leu Arg Thr Pro Ala Gly Gly Lys Gly Leu Ala Leu Ser Ser Ser Lys Val Lys Ser Val Val Met Val Val Phe Asp 280 Asn Asp Lys Val Pro Val Glu Gln Leu Arg Phe Trp Arg His Trp His Ser Arg Gln Pro Thr Ala Lys Gln Arg Val Ile Asp Val Ala Asp Cys Lys Glu Asn Phe Asn Thr Val Gln His Ile Glu Glu Val Ala Tyr Asn Ala Leu Ser Phe Val Trp Asn Val Asn Glu Glu Ala Lys Val Phe Ile Gly Val Asn Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Val Lys Gly Val Pro Leu Asn Leu Gln Ile Asp Thr Tyr Asp Cys Gly Ala Gly Thr Glu Arg Leu Val His Arg Ala Val Cys Gln Ile Lys Ile Phe Cys 390 Asp Lys Gly Ala Glu Arg Lys Met Arg Asp Asp Glu Arg Lys Gln Phe Arg Arg Lys Val Lys Cys Pro Asp Ser Ser Asn Asn Ala Gly Ile Lys Gly Cys Leu Leu Ser Gly Phe Arg Gly Asn Glu Thr Thr Tyr Leu Arg Pro Glu Thr Asp Leu Glu Thr Gln Pro Val Leu Phe Ile Pro Asn Leu 455 His Phe Ser Ser Leu Gln Arg Pro Gly Gly Val Val Pro Ser Ala Gly 470 His Ser Ser Ser Asp Arg Leu Pro Leu Lys Arg Thr Cys Ser Pro Phe 490 485 Ala Glu Glu Phe Glu Pro Leu Pro Ser Lys Gln Ala Lys Glu Asp Asp Leu Gln Arg Val Leu Leu Tyr Val Arg Arg Glu Thr Glu Glu Val Phe Asp Ala Leu Met Leu Lys Thr Pro Asp Leu Lys Gly Leu Arg Asn Ala Ile Ser Glu Lys Tyr Gly Leu Pro Glu Glu Asn Ile Cys Lys Val Tyr 545 Lys Lys Cys Lys Arg Gly Ile Leu Val Asn Met Asp Asn Asn Ile Ile Gln His Tyr Ser Asn His Val Ala Phe Leu Leu Asp Met Gly Glu Leu Asp Gly Lys Ile Gln Ile Ile Leu Lys Glu Leu <210> 17 4840 <211> DNA <213> Drosophila <400> 17 aaaaatagaa aaaacaacaa caaattggct tgaaaacgca aatgccaggc gcaacgcccc 60 cgaaccgacc cgcccctca acttttgcgc cctccagtag caatagcagc aatatgagca 120 gcagcaacat caaatgttag gccaaaatgc acaaaccgcc agcaacaaag gcagcaccaa 180 gcgaacgaaa caacaacagc tccacatacc acaaagagtg gcacattaga agcggccaaa 240 agcagccagc cgagagcatt gtgtaagcca aaggcccaga gagccaggct aaaagccccc 300 360 agacgcacaa caacaacaac aacaactaaa acagcacaaa gagtggcgaa aggtgcaccc accagcaaaa cagcaacaac ggagcaacca acaacagcag cagcagcagc agcagccaca 420 480 tttcagttac agctccagac tcccaggttg cagactccca aagcaaacag actccagtcc acgatecage tecagiteca ecgatecgat ceaetgetee agegigeteg agigecatag 540 600 atoctoacca agtgocaaaa toogoatoot gatoocaaga gotoaaggoa coooggocoa aaattgagct gagaacgaaa cgaaggaagt tccttagtgc catagaaagc agttaatgaa 660 acaacgacta agacgaagat cgaccatcca gaaccggagg gagctaattg cgaacgaaag 720 aaaccacaaa gtgccttcca tcaatccgtt gataagtgat atttattatg tttatacttg 780 ccagcagceg aggcagcaac agcaatagca acaaccatag gggatcacgg catcgatgat 840 cagtccacga ccaagtccta gtgcaatccg gaatccagtt caaattagtt caataagccg 900 tatetaceae gtataatgte cacateeace gecacaaega gegttateae gtecaaegag 960 1020 ctctcgctgt ccggccacgc ccacggtcac ggtcacgccc accagttgca ccagcacacc cacagoogoc taggagttgg cgttggtgtt ggcatcetta gcgacgcatc cctatcgccc 1080

atccaacaag gcagtggcgg ccacagcggc ggaggtaaca caaacagttc accactggcg .

cccaacggag tgccacttct cacaacaatg caccgatcac cggactcacc gcagccagaa 1200 ttggccacca tgacgaacgt caacgtgctg gatctgcaca cggataactc caagctgtac 1260 gacaaggagg ctgtatttat atacgaaacg cccaaggtgg tgatgccagc ggatggcggg 1320 ggtggcaata attccgatga aggtcatgcc atcgatgcgc ggattgcggc ccaaatgggc 1380 aaccaagccc agcaacagca gcagcagcaa cagcagacgg aacaccagcc gctggccaag 1440 atcgagttcg atgagaacca gataatccgg gtggtgggac caaatggcga gcaacagcaa 1500 atcatetege gggagateat caatggggag cateatatee tgtegegaaa egaggetggt 1560 gagcacattc tcacacggat cgtcagtgat ccctccaagt tgatgcccaa tgacaatgca 1620 gtggccacgg ccatgtacaa ccaggcccaa aagatgaaca atgatcacgg gcaggcggta 1680 tatcagacat caccattgcc gctagacgcg tctgtattgc attatagtgg cggcaatgat 1740 togaatgtaa ttaagaogga ggoogatato taogaggato acaagaaaca tgoggotgca 1800 gcagcagctg ctgccggcgg aggatccatc atatacacca catccgatcc gaacggagtg 1860 aatgtgaaac aactgcccca tttgacggta ccccaaaaac ttgatcccga cctctatcaa 1920 geogataage atatagattt gatetacaac gatggcagea agacggtgat ttactccact 1980 acggatcaga agagtttgga aatatactcg ggcggcgaca tcggcagcct ggtgtccgac 2040 ggccaagtgg tggtccaggc gggactgccg tatgccacca ccaccggagc cggcggccag 2100 cccgtctata tcgtggccga cggtgccttg ccagcgggag tcgaggagca tctgcagagt 2160 ggaaagetea atggeeagae cacacetate gatgtetetg geetategea aaatgagatt 2220 caaggetttt tgeteggete acaeceeteg teateggega eggtaageac aaceggegtt 2280 gtetecacga caacgatete geateaceag caacageage ageageagea acageaacag 2340 cagcagcagc agcagcaaca ccagcagcag cagcaacatc ccggcgacat tgttagtgcc 2400 gctggcgtgg ggagcacggg ctccattgtc tcctctgcgg cgcaacagca gcagcagcag 2460 caactaatta gcatcaaacg agagcccgaa gacttgcgca aggatcccaa gaatggcaac 2520 attgccggtg cagcaacagc aaatggaccc ggttcggtca taacccaaaa gtcctttgat 2580 tatacggaat tgtgccagcc gggcacgctg atcgatgcca atggcagcat acccgtcagc 2640 gtgaacagca tccagcagag aacggcggtc catggcagcc agaacagtcc caccacatcg 2700 ctggtggaca' ccagcaccaa tggatccacg cgatcgcggc cctggcacga ctttggacgt 2760 cagaatgatg cegacaaaat acaaatacca aaaatettea caaacgtggg etteegatat 2820

cacctggaga	gccccatcag	ttcatcgcag	aggcgcgagg	acgatcgcat	cacctacatc	2880
aacaagggtc	aattctatgg	aataacgctg	gagtatgtgc	acgatgcgga	aaagcccatt	2940
aagaacacca	ccgtcaagag	tgtgatcatg	ctaatgttcc	gcgaggagaa	gagtcccgag	3000
gatgaġatca	aggcctggca	attctggcac	agtegteage	attccgtgaa	gcagagaatc	3060
ttggatgcag	atacgaagaa	ctcggttggc	ctcgttggct	gcatcgagga	agtgtcgcac	3120
aatgccatcg	ccgtctactg	gaatccgctg	gagageteeg	ccaagatcaa	cattgcggtt	3180
cagtgcttga	gcacggattt	cagcagtcaa	aagggaggcc	tgccgctgca	cgtacaaatc	3240
gacacatttg	aggaccccag	agatacggcg	gtcttccacc	gcggctactg	tcagataaag	3300
gtcttctgcg	ataagggcgc	cgaacgaaag	acgcgcgatg	aagagcggcg	ggccgccaaa	3360
cgaaagatga	cagccacggg	cagaaagaag	ctggacgagc	tttaccatcc	ggtaacggat	3420
cggtccgagt	tctatggcat	gcaggacttc	gccaagccgc	cggtgctatt	ctegeeegee	3480
gaggacatgg	agaaggtagg	tcagctgggc	attggcgctg	ccaccggcat	gacattcaac	3540
cccctgagca	acggcaactc	caactccaac	tcgcactcgt	ccttgcagag	cttctacggc	3600
catgagactg	actcgccgga	cctgaagggg	gcctcaccgt	tcctgctcca	cggccagaag	3660
gtggccacge	cgacgeteaa	gttccacaac	cattttccgc	ccgacatgca	gaccgataag	3720
aaggatcaca	tactggacca	gaacatgttg	accagcacac	ccctgaccga	ctttggtccg	3780
ccgatgaagc	gcggcaggat	gacgccgccg	acctcggaac	gcgtgatgct	gtacgtgcgg	3840
caggagaacg	aggaggtgta	tacaccgttg	cacgtggtgc	cgcccaccac	gatcggcctg	3900
ctaaatgcga	ttgaaaacaa	atacaaaatc	tcaacaacga	gcataaataa	catttatcgc	3960
acaaacaaga	aggggattac	tgcgaaaatt	gacgatgaca	tgatatcgtt	ctactgcaac	4020
gaggacatct	ttctgctgga	ggtgcaacag	atcgaggacg	acctgtacga	tgtgacgctc	4080
acggagetge	ccaatcagta	gcgctggcag	tacgggtagc	accegetaac	cgcactcaaa	4140
aaaaaagca	aacaaacaca	caaattacgg	acacaacaag	ttgtttcaat	aagccatttt	4200
ccatagagcc	taagtetaaa	tatcgtagtt	ataataatgg	gateegeaac	aaatcgagtt	4260
gcaacgaatg	ttaagaacgc	taacacaata	cgcatgtaaa	atgatacttt	aaaattgatt	4320
tagttatttt	agcaacaatg	agattatcta	aaattgtttg	atcaaatttt	acattctcgc	4380
tatgtctata	gataattcta	agcccgtaag	cccataagcg	taatcgtaat	cgtaatcgta	4440
ccgtgtattt	atgctcatat	ataaacaact	atatatatat	atatatatat	atatatgtgc	4500

ggagtgcaac agtgtctgtc cagtaggaga taagtctcgt ttccgc	tccc ctgcttatgc 4560
tatgacetta ggtecaggge aagtatgagt tacegaatet atetat	tagg tgcatctaac 4620
gaaaggaate attagetetg cacgaactet ageegtagee tattgt	aatc catttgtatg 4680
tttggcttaa gcgttttact tgttgaatat aaagtgtaaa attatt	tttg aaaaaaaaaa 4740
acccacacaa aacacaaatc gtttgttcta tatttctgtt tcaaaa	actaa ctcgttaccc 4800
acaatcccct ctgttatgta taattaggat ctctgtacac	4840
.010. 10	

<210> 18

<211> 1061

<212> PRT

<213> murine

<400> 18

Met Ser Thr Ser Thr Ala Thr Thr Ser Val Ile Thr Ser Asn Glu Leu

1 5 . 10 15

Ser Leu Ser Gly His Ala His Gly His Gly His Ala His Gln Leu His 20 25 30

Gln His Thr His Ser Arg Leu Gly Val Gly Val Gly Val Gly Ile Leu
35 40 45

Ser Asp Ala Ser Leu Ser Pro Ile Gln Gln Gly Ser Gly Gly His Ser 50 55 60

Gly Gly Asn Thr Asn Ser Ser Pro Leu Ala Pro Asn Gly Val Pro 65 70 75 80

Leu Leu Thr Thr Met His Arg Ser Pro Asp Ser Pro Gln Pro Glu Leu 85 90 95

Ala Thr Met Thr Asn Val Asn Val Leu Asp Leu His Thr Asp Asn Ser 100 105 110

Lys Leu Tyr Asp Lys Glu Ala Val Phe Ile Tyr Glu Thr Pro Lys Val 115 120 125

Val Met Pro Ala Asp Gly Gly Gly Gly Asn Asn Ser Asp Glu Gly His 130 135 140

Ala Ile Asp Ala Arg Ile Ala Ala Gln Met Gly Asn Gln Ala Gln Gln 145 150 155 160

Gln Gln Gln Gln Gln Gln Thr Glu His Gln Pro Leu Ala Lys Ile 165 170 175

Glu Phe Asp Glu Asn Gln Ile Ile Arg Val Val Gly Pro Asn Gly Glu 180 185 190

Gln Gln Gln Ile Ile Ser Arg Glu Ile Ile Asn Gly Glu His His Ile

195 200 205 Leu Ser Arg Asn Glu Ala Gly Glu His Ile Leu Thr Arg Ile Val Ser Asp Pro Ser Lys Leu Met Pro Asn Asp Asn Ala Val Ala Thr Ala Met 230 Tyr Asn Gln Ala Gln Lys Met Asn Asn Asp His Gly Gln Ala Val Tyr Gln Thr Ser Pro Leu Pro Leu Asp Ala Ser Val Leu His Tyr Ser Gly Gly Asn Asp Ser Asn Val Ile Lys Thr Glu Ala Asp Ile Tyr Glu Asp 280 His Lys Lys His Ala Ala Ala Ala Ala Ala Ala Gly Gly Gly Ser Ile Ile Tyr Thr Thr Ser Asp Pro Asn Gly Val Asn Val Lys Gln Leu 310 315 Pro His Leu Thr Val Pro Gln Lys Leu Asp Pro Asp Leu Tyr Gln Ala Asp Lys His Ile Asp Leu Ile Tyr Asn Asp Gly Ser Lys Thr Val Ile Tyr Ser Thr Thr Asp Gln Lys Ser Leu Glu Ile Tyr Ser Gly Gly Asp Ile Gly Ser Leu Val Ser Asp Gly Gln Val Val Val Gln Ala Gly Leu Pro Tyr Ala Thr Thr Gly Ala Gly Gly Gln Pro Val Tyr Ile Val Ala Asp Gly Ala Leu Pro Ala Gly Val Glu Glu His Leu Gln Ser Gly Lys Leu Asn Gly Gln Thr Thr Pro Ile Asp Val Ser Gly Leu Ser Gln 420 425 Asn Glu Ile Gln Gly Phe Leu Leu Gly Ser His Pro Ser Ser Ser Ala Thr Val Ser Thr Thr Gly Val Val Ser Thr Thr Thr Ile Ser His His 450 470 475 Gln His Gln Gln Gln Gln His Pro Gly Asp Ile Val Ser Ala Ala 485

Gly Val Gly Ser Thr Gly Ser Ile Val Ser Ser Ala Ala Gln Gln 505 Gln Gln Gln Leu Ile Ser Ile Lys Arg Glu Pro Glu Asp Leu Arg Lys Asp Pro Lys Asn Gly Asn Ile Ala Gly Ala Ala Thr Ala Asn Gly 530 Pro Gly Ser Val Ile Thr Gln Lys Ser Phe Asp Tyr Thr Glu Leu Cys 555 Gln Pro Gly Thr Leu Ile Asp Ala Asn Gly Ser Ile Pro Val Ser Val Asn Ser Ile Gln Gln Arg Thr Ala Val His Gly Ser Gln Asn Ser Pro Thr Thr Ser Leu Val Asp Thr Ser Thr Asn Gly Ser Thr Arg Ser Arg Pro Trp His Asp Phe Gly Arg Gln Asn Asp Ala Asp Lys Ile Gln Ile 615 620 Pro Lys Ile Phe Thr Asn Val Gly Phe Arg Tyr His Leu Glu Ser Pro 630 Ile Ser Ser Ser Gln Arg Arg Glu Asp Asp Arg Ile Thr Tyr Ile Asn 650 Lys Gly Gln Phe Tyr Gly Ile Thr Leu Glu Tyr Val His Asp Ala Glu 665 Lys Pro Ile Lys Asn Thr Thr Val Lys Ser Val Ile Met Leu Met Phe Arg Glu Glu Lys Ser Pro Glu Asp Glu Ile Lys Ala Trp Gln Phe Trp 690 695 His Ser Arg Gln His Ser Val Lys Gln Arg Ile Leu Asp Ala Asp Thr Lys Asn Ser Val Gly Leu Val Gly Cys Ile Glu Glu Val Ser His Asn 725 Ala Ile Ala Val Tyr Trp Asn Pro Leu Glu Ser Ser Ala Lys Ile Asn 745 Ile Ala Val Gln Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Gly 760 Leu Pro Leu His Val Gln Ile Asp Thr Phe Glu Asp Pro Arg Asp Thr 775 Ala Val Phe His Arg Gly Tyr Cys Gln Ile Lys Val Phe Cys Asp Lys 790

Gly Ala Glu Arg Lys Thr Arg Asp Glu Glu Arg Arg Ala Ala Lys Arg 805 810 815

Lys Met Thr Ala Thr Gly Arg Lys Leu Asp Glu Leu Tyr His Pro 820 825 830

Val Thr Asp Arg Ser Glu Phe Tyr Gly Met Gln Asp Phe Ala Lys Pro 835 840 845

Pro Val Leu Phe Ser Pro Ala Glu Asp Met Glu Lys Val Gly Gln Leu 850 855 860

Gly Ile Gly Ala Ala Thr Gly Met Thr Phe Asn Pro Leu Ser Asn Gly 865 870 875 880

Asn Ser Asn Ser Asn Ser His Ser Ser Leu Gln Ser Phe Tyr Gly His 885 890 895

Glu Thr Asp Ser Pro Asp Leu Lys Gly Ala Ser Pro Phe Leu Leu His 900 905 910

Gly Gln Lys Val Ala Thr Pro Thr Leu Lys Phe His Asn His Phe Pro 915 920 925

Pro Asp Met Gln Thr Asp Lys Lys Asp His Ile Leu Asp Gln Asn Met 930 935 940

Leu Thr Ser Thr Pro Leu Thr Asp Phe Gly Pro Pro Met Lys Arg Gly 945 950 955 960

Arg Met Thr Pro Pro Thr Ser Glu Arg Val Met Leu Tyr Val Arg Gln
965 970 975

Glu Asn Glu Glu Val Tyr Thr Pro Leu His Val Val Pro Pro Thr Thr 980 985 990

Ile Gly Leu Leu Asn Ala Ile Glu Asn Lys Tyr Lys Ile Ser Thr Thr 995 1000 1005

Ser Ile Asn Asn Ile Tyr Arg Thr Asn Lys Lys Gly Ile Thr Ala 1010 1015 1020

Lys Ile Asp Asp Asp Met Ile Ser Phe Tyr Cys Asn Glu Asp Ile 1025 1030 1035

Phe Leu Leu Glu Val Gln Gln Ile Glu Asp Asp Leu Tyr Asp Val 1040 1045 1050

Thr Leu Thr Glu Leu Pro Asn Gln 1055 1060

<210> 19

<211> 21

<212> DNA

<213> artificial sequence

<220> <223> human p49 mgr <400> 19 gaagtetttg atgeeetgat g 21 <210> 20 <211> 21 <212> DNA <213> artificial sequence <220> <223> human p49 mgr <400> 20 'aacccattcc ctcgacatag a 21 <210> 21 <211> 20 ( <212> DNA <213> artificial sequence <220> <223> human p70 mgr <400> 21 agcgcgatga cacaggagta 20 <210> 22 <211> 20 <212> DNA <213> artificial sequence <220> <223> human p70 mgr <400> 22 20 cgttgctatg gagacagtga <210> 23 <211> 20 <212> DNA <213> artificial sequence <220> <223> human bom <400> 23 ccgtttaaca aggacactgc 20 <210> 24 <211> 20 <212> DNA <213> artificial sequence

. .

<220>		
<223>	human bom	
<400>	24	
	gcca ccaaatctct	20
organi	300a 00aaa0000	
<210>	25	
<211>	20	
<212>	DNA	
<213>	artificial sequence	
<220>		
<223>	murine p70 mgr	
-400-	25	
<400>	25 atga cacaggagta	20
agegeg	acya cacaggagca	
<210>	26	
<211>		
<212>		
<213>	artificial sequence	
		•
<220>		
<223>	murine p70 mgr	
-400-	26	
<400>	26 agag ctgaactgat	20
agegee	agag cogaacegac	
<210>	27	
<211>	20	
<212>		
<213>	artificial sequence	
<220>		
<223>	murine p61 mgr	
<400>	27	
	aggtt cettgagtte	20
	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
<210>	28	
<211>	20	
<212>		
<213>	artificial sequence	
<220>	murine p61 mgr	
	murine por mgr	
<400>	28	
	cagag ctgaactgat	20
J		
<210>		
<211>		
<212>		
<213>	artificial sequence	
<220>		
5440>		

<213> Drosophila

<223> murine bom <400> 29 aaaggggagc gagttcattg 20 <210> 30 <211> 20 <212> DNA <213> artificial sequence <220> <223> murine bom <400> 30 20 agageteteg gtgatggata <210> 31 . <211> 34 <212> DNA <213> artificial sequence <223> Drosophila dopa decarboxylase promoter <400> 31 ggtggtgctc taataaccgg tttccaagat gcgc 34 <210> 32 <211> 34 <212> DNA <213> artificial sequence <220> <223> Drosophila PCNA promoter gggtaaaaag tgtgaacaat caaaccagtt ggca 34 <210> 33 <211> 84 <212> DNA <213> artificial sequence. <220> <223> Human engrailed-1 promoter <400> 33 ggacacacac ccaaacccac acccaccac aaacacacaa accggcagtg acaacaacca 84 cccatccttc aataacagca acca <210> 34 <211> 4747 <212> DNA

<400> 34 aaaaatagaa aaaacaacaa caaattggct tgaaaacgca aatgccaggc gcaacgcccc 60 cgaaccgacc egececetea aettttgege eetecagtag caatageage aatatgagea 120 gcagcaacat caaatgttag gccaaaatgc acaaaccgcc agcaacaaag gcagcacaa 180 gcgaacgaaa caacaacagc tccacatacc acaaagagtg gcacattaga agcggccaaa 240 agcagecage egagageatt gtgtaageca aaggeecaga gagecagget aaaageecee 300 agacgcacaa caacaacaac aacaactaaa acagcacaaa gagtggcgaa aggtgcaccc 360 accagcasaa cagcaacaac ggagcaacca acaacagcag cagcagcagc agcagccaca 420 tttcagttac agetecagac teccaggttg cagactecca aageaaacag actecagtee 480 acgatecage tecagities cogatecgat coactgetee agegigeteg agitgecatag 540 atcoteacea agtgecaaaa teegoateet gateecaaga geteaaggea ceeeggeeca 600 aaattgaget gagaacgaaa egaaggaagt teettagtge catagaaage agttaatgaa 660 acaacgacta agacgaagat cgaccatcca gaaccggagg gagctaattg cgaacgaaag 720 aaaccacaaa gtgccttcca tcaatccgtt gataagtgat atttattatg tttatacttg 780 ccagcagccg aggcagcaac agcaatagca acaaccatag gggatcacgg catcgatgat 840 cagtccacga ccaagtccta gtgcaatccg gaatccagtt caaattagtt caataagccg 900 tatctaccac gtataatgtc cacatccacc gccacaacga gcgttatcac gtccaacgag 960 ctctcgctgt ccggccacgc ccacggtcac ggtcacgccc accagttgca ccagcacacc 1020 cacagoogoo taggagttgg ogttggtgtt ggcatootta gcgacgcato cotatogooc 1080 atccaacaag gcagtggcgg ccacagcggc ggaggtaaca caaacagttc accactggcg 1140 eccaacggag tgccacttct cacaacaatg caccgatcac cggactcacc gcagccagaa 1200 ttggccacca tgacgaacgt caacgtgctg gatctgcaca cggataactc caagctgtac 1260 gacaaggagg ctgtatttat atacgaaacg cccaaggtgg tgatgccagc ggatggcggg 1320 ggtggcaata attccgatga aggtcatgcc atcgatgcgc ggattgcggc ccaaatgggc 1380 aaccaagccc agcaacagca gcagcagcaa cagcagacgg aacaccagcc gctggccaag 1440 atcgagttcg atgagaacca gataatccgg gtggtgggac caaatggcga gcaacagcaa 1500 atcatctcgc gggagatcat caatggggag catcatatcc tgtcgcgaaa cgaggctggt 1560 gagcacatte teacaeggat egteagtgat ecetecaagt tgatgeecaa tgacaatgea 1620

gtggccacgg ccatgtacaa ccaggcccaa aagatgaaca atgatcacgg gcaggcggta 1680 tatcagacat caccattgcc gctagacgcg tctgtattgc attatagtgg cggcaatgat 1740 tcgaatgtaa ttaagacgga ggccgatatc tacgaggatc acaagaaaca tgcggctgca 1800 gcagcagctg ctgccggcgg aggatccatc atatacacca catccgatcc gaacggagtg 1860 aatgtgaaac aactgcccca tttgacggta ccccaaaaac ttgatcccga cctctatcaa 1920 gccgataagc atatagattt gatctacaac gatggcagca agacggtgat ttactccact 1980 2040 acqqatcaga agagtttgga aatatactcg ggcggcgaca tcggcagcct ggtgtccgac ggccaagtgg tggtccaggc gggactgccg tatgccacca ccaccggagc cggcggccag 2100 2160 cccgtctata tcgtggccga cggtgccttg ccagcgggag tcgaggagca tctgcagagt ggaaagetea atggeeagae cacacetate gatgtetetg geetategea aaatgagatt 2220 2280 caaggetttt tgeteggete acacceteg teateggega eggtaageac aaceggegtt gtotocacga caacgatoto gcatcaccag caacagcagc agcagcagca acagcaacag 2340 cagcagcagc agcagcaaca ccagcagcag cagcaacatc ccggcgacat tgttagtgcc 2400 2460 getggegtgg ggageaeggg etecattgte teetetgegg egeaaeagea geageageag 2520 caactaatta gcatcaaacg agagcccgaa gacttgcgca aggatcccaa gaatggcaac 2580 attgccggtg cagcaacagc aaatggaccc ggttcggtca taacccaaaa gtcctttgat tatacggaat tgtgccagcc gggcacgctg atcgatgcca atggcagcat acccgtcagc 2640 2700 gtgaacagca tccagcagag aacggcggtc catggcagcc agaacagtcc caccacatcg ctggtggaca ccagcaccaa tggatccacg cgatcgcggc cctggcacga ctttggacgt 2760 2820 cagaatgatg ccgacaaaat acaaatacca aaaatcttca caaacgtggg cttccgatat 2880 cacctggaga gccccatcag ttcatcgcag aggcgcgagg acgatcgcat cacctacatc 2940 aacaagggtc aattctatgg aataacgctg gagtatgtgc acgatgcgga aaagcccatt 3000 aagaacacca ccgtcaagag tgtgatcatg ctaatgttcc gcgaggagaa gagtcccgag 3060 gatgagatca aggcctggca attctggcac agtcgtcagc attccgtgaa gcagagaatc 3120 ttggatgcag atacgaagaa eteggttgge etegttgget geategagga agtgtegeae 3180 aatgccatcg ccgtctactg gaatccgctg gagagctccg ccaagatcaa cattgcggtt 3240 cagtgcttga gcacggattt cagcagtcaa aagggaggcc tgccgctgca cgtacaaatc gacacatttg aggaccccag agatacggcg gtcttccacc gcggctactg tcagataaag 3300

gtcttctgcg	ataagggcgc	cgaacgaaag	acgcgcgatg	aagagcggcg	ggccgccaaa	3360
cgaaagatga	cagccacggg	çagaaagaag	ctggacgagc	tttaccatcc	ggtaacggat	3420
cggtccgagt	tctatggcat	gcaggacttc	gccaagccgc	cggtgctatt	ctcgcccgcc	3480
gaggacatgg	agaagagctt	ctacggccat	gagactgact	cgccggacct	gaagggggcc	3540
tcaccgttcc	tgctccacgg	ccagaaggtg	gccacgccga	cgctcaagtt	ccacaaccat	3600
tttccgcccg	acatgcagac	cgataagaag	gatcacatac	tggaccagaa	catgttgacc	3660
agçacacccc	tgaccgactt	tggtccgccg	atgaagcgcg	gcaggatgac	gccgccgacc	3720
tcggaacgcg	tgatgctgta	cgtgcggcag	gagaacgagg	aggtgtatac	accgttgcac	3780
gtggtgccgc	ccaccacgat	cggcctgcta	aatgcgattg	aaaacaaata	caaaatctca	3840
acaacgagca	taaataacat	ttatcgcaca	aacaagaagg	ggattactgc	gaaaattgac	3900
gatgacatga	tatcgttcta	ctgcaacgag	gacatettte	tgctggaggt	gcaacagatc	`3960
gaggacgacc	tgtacgatgt	gacgeteacg	gagetgeeca	atcagtagcg	ctggcagtac	4020
gggtagcacc	cgctaaccgc	actcaaaaaa	aaaagcaaac	aaacacacaa	attacggaca	4080
caacaagttg	tttcaataag	ccattttcca	tagageetaa	gtctaaatat	cgtagttata	4140
ataatgggat	ccgcaacaaa	tcgagttgca	acgaatgtta	agaacgctaa	cacaatacgc	4200
atgtaaaatg	atactttaaa	attgatttag	ttattttagc	aacaatgaga	ttatctaaaa	42.60
ttgtttgatc	aaattttaca	ttctcgctat	gtctatagat	aattctaagc	ccgtaagccc	4320
ataagcgtaa	tcgtaatcgt	aatcgtaccg	tgtatttatg	ctcatatata	aacaactata	4380
tatatatata	tatatatata	tatgtgcgga	gtgcaacagt	gtctgtccag	taggagataa	4440
gtctcgtttc	cgctcccctg	cttatgctat	gaccttaggt	ccagggcaag	tatgagttac	4500
cgaatctatc	tattaggtgc	atctaacgaa	aggaatcatt	agctctgcac	gaactctagc	4560
cgtagcctat	tgtaatccat	ttgtatgttt	ggcttaagcg	ttttacttgt	tgaatataaa	4620
gtgtaaaatt	atttttgaaa	aaaaaaaacc	cacacaaaac	acaaatcgtt	tgttctatat	4680
ttctgtttca	aaactaactc	gttacccaca	atcccctctg	ttatgtataa	ttaggatctc	4740
tgtacac						4747

<sup>&</sup>lt;210> 35

<sup>&</sup>lt;211> 1030

<sup>&</sup>lt;212> PRT

<sup>&</sup>lt;213> Drosophila

<400> 35 Met Ser Thr Ser Thr Ala Thr Thr Ser Val Ile Thr Ser Asn Glu Leu Ser Leu Ser Gly His Ala His Gly His Gly His Ala His Gln Leu His Gln His Thr His Ser Arg Leu Gly Val Gly Val Gly Val Gly Ile Leu Ser Asp Ala Ser Leu Ser Pro Ile Gln Gln Gly Ser Gly Gly His Ser Gly Gly Gly Asn Thr Asn Ser Ser Pro Leu Ala Pro Asn Gly Val Pro Leu Leu Thr Thr Met His Arg Ser Pro Asp Ser Pro Gln Pro Glu Leu 90 Ala Thr Met Thr Asn Val Asn Val Leu Asp Leu His Thr Asp Asn Ser Lys Leu Tyr Asp Lys Glu Ala Val Phe Ile Tyr Glu Thr Pro Lys Val Val Met Pro Ala Asp Gly Gly Gly Asn Asn Ser Asp Glu Gly His 135 Ala Ile Asp Ala Arg Ile Ala Ala Gln Met Gly Asn Gln Ala Gln Gln Gln Gln Gln Gln Gln Gln Thr Glu His Gln Pro Leu Ala Lys Ile Glu Phe Asp Glu Asn Gln Ile Ile Arg Val Val Gly Pro Asn Gly Glu 185 Gln Gln Gln Ile Ile Ser Arg Glu Ile Ile Asn Gly Glu His His Ile 205 Leu Ser Arg Asn Glu Ala Gly Glu His Ile Leu Thr Arg Ile Val Ser 220 215 Asp Pro Ser Lys Leu Met Pro Asn Asp Asn Ala Val Ala Thr Ala Met 235 Tyr Asn Gln Ala Gln Lys Met Asn Asn Asp His Gly Gln Ala Val Tyr Gln Thr Ser Pro Leu Pro Leu Asp Ala Ser Val Leu His Tyr Ser Gly Gly Asn Asp Ser Asn Val Ile Lys Thr Glu Ala Asp Ile Tyr Glu Asp His Lys Lys His Ala Ala Ala Ala Ala Ala Ala Ala Gly Gly Ser

290 295 300 Ile Ile Tyr Thr Thr Ser Asp Pro Asn Gly Val Asn Val Lys Gln Leu Pro His Leu Thr Val Pro Gln Lys Leu Asp Pro Asp Leu Tyr Gln Ala 330 Asp Lys His Ile Asp Leu Ile Tyr Asn Asp Gly Ser Lys Thr Val Ile Tyr Ser Thr Thr Asp Gln Lys Ser Leu Glu Ile Tyr Ser Gly Gly Asp Ile Gly Ser Leu Val Ser Asp Gly Gln Val Val Val Gln Ala Gly Leu Pro Tyr Ala Thr Thr Gly Ala Gly Gly Gln Pro Val Tyr Ile Val 390 Ala Asp Gly Ala Leu Pro Ala Gly Val Glu Glu His Leu Gln Ser Gly Lys Leu Asn Gly Gln Thr Thr Pro Ile Asp Val Ser Gly Leu Ser Gln Asn Glu Ile Gln Gly Phe Leu Leu Gly Ser His Pro Ser Ser Ser Ala 440 Thr Val Ser Thr Thr Gly Val Val Ser Thr Thr Thr Ile Ser His His 470 Gln His Gln Gln Gln Gln His Pro Gly Asp Ile Val Ser Ala Ala 490 Gly Val Gly Ser Thr Gly Ser Ile Val Ser Ser Ala Ala Gln Gln Gln Gln Gln Leu Ile Ser Ile Lys Arg Glu Pro Glu Asp Leu Arg 520 Lys Asp Pro Lys Asn Gly Asn Ile Ala Gly Ala Ala Thr Ala Asn Gly Pro Gly Ser Val Ile Thr Gln Lys Ser Phe Asp Tyr Thr Glu Leu Cys 550 Gln Pro Gly Thr Leu Ile Asp Ala Asn Gly Ser Ile Pro Val Ser Val Asn Ser Ile Gln Gln Arg Thr Ala Val His Gly Ser Gln Asn Ser Pro

Thr Thr Ser Leu Val Asp Thr Ser Thr Asn Gly Ser Thr Arg Ser Arg 595 600 Pro Trp His Asp Phe Gly Arg Gln Asn Asp Ala Asp Lys Ile Gln Ile 615 Pro Lys Ile Phe Thr Asn Val Gly Phe Arg Tyr His Leu Glu Ser Pro Ile Ser Ser Ser Gln Arg Arg Glu Asp Asp Arg Ile Thr Tyr Ile Asn Lys Gly Gln Phe Tyr Gly Ile Thr Leu Glu Tyr Val His Asp Ala Glu Lys Pro Ile Lys Asn Thr Thr Val Lys Ser Val Ile Met Leu Met Phe 680 Arg Glu Glu Lys Ser Pro Glu Asp Glu Ile Lys Ala Trp Gln Phe Trp 695 His Ser Arg Gln His Ser Val Lys Gln Arg Ile Leu Asp Ala Asp Thr Lys Asn Ser Val Gly Leu Val Gly Cys Ile Glu Glu Val Ser His Asn Ala Ile Ala Val Tyr Trp Asn Pro Leu Glu Ser Ser Ala Lys Ile Asn 745 Ile Ala Val Gln Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Gly Leu Pro Leu His Val Gln Ile Asp Thr Phe Glu Asp Pro Arg Asp Thr Ala Val Phe His Arg Gly Tyr Cys Gln Ile Lys Val Phe Cys Asp Lys Gly Ala Glu Arg Lys Thr Arg Asp Glu Glu Arg Arg Ala Ala Lys Arg 805 Lys Met Thr Ala Thr Gly Arg Lys Leu Asp Glu Leu Tyr His Pro Val Thr Asp Arg Ser Glu Phe Tyr Gly Met Gln Asp Phe Ala Lys Pro Pro Val Leu Phe Ser Pro Ala Glu Asp Met Glu Lys Ser Phe Tyr Gly 855 His Glu Thr Asp Ser Pro Asp Leu Lys Gly Ala Ser Pro Phe Leu Leu His Gly Gln Lys Val Ala Thr Pro Thr Leu Lys Phe His Asn His Phe

Pro	Pro	Asp	Met 900	Gln	Thr	Asp	Гув	Lys 905	Asp	His	Ile	Leu	Asp 910	Gln	Asn	
Met	Leu	Thr 915	Ser	Thr	Pro	Leu	Thr 920	qaA	Phe	Gly	Pro	Pro 925	Met	Lys	Arg	
Gly	Arg 930	Met	Thr	Pro	Pro	Thr 935	Ser	Glu	Arg	Val	Met 940	Leu	Tyr	Val	Arg	
Gln 945	Glu	Asn	Glu	Glu	Val 950	Tyr	Thr	Pro	Leu	His 955	Va1	Val	Pro	Pro	Thr 960	
Thr	Ile	Gly	Leu	Leu 965	Asn	Ala	Ile	Glu	Asn 970	Lys	Tyr	Lys	Ile	Ser 975	Thr	
Thr	Ser	Ile	Asn 980	Asn	Ile	Tyr	Arg	Thr 985	Asn	ГÀв	Lys	Gly	Ile 990	Thr	Ala	
Lys	Ile	Asp 995	Asp	Asp	Met	Ile	Ser 100		е Ту:	r Cýs	a A's:	n Gl		sp I	ļe Phe	
Leu	Leu 101		u Va	1 Gl:	n Gl	n Il		lu A	sp A	sp Le		yr 020	Asp	Val '	Thr	
Leu	Thr 102		u Le	u Pr	o As	n Gl:										
<21 <21 <21	1>	36 5650 DNA														
<21			ophi	la												
<40 aaa		36 gaa	aaaa	.caac	aa c	aaat	tggc	t tg	aaaa	.cgca	aat	gcca	ggc	gcaa	.cgcccc	60
cga	accg	acc	cgcc	ccct	ca a	cttt	tgcg	c cc	tcca	gtag	caa	tago	agc	aata	tgagca	120
gca	gcaa	cat	caaa	tgtt	ag g	ccaa	aatg	c ac	aaac	cgcc	ago	aaca	aag	gcag	caccaa	180
gcg	aacg	aaa	caac	aaca	gc t	ccac	atac	c ac	aaag	jagtg	gca	catt	aga	agcg	gccaaa	240
ago	agcó	agc	cgag	gagca	itt g	jtgta	agco	a aa	ıggco	caga	gag	gccaç	gct	aaaa	gcccc	300
aga	cgca	caa	caac	aaca	ac a	acaa	ctae	a ac	agca	caaa	gaç	gtggd	gaa	aggt	gcaccc	360
acc	agca	ıaaa	cago	caaca	ac g	gago	aaco	ca ac	caaca	agcag	cag	gcago	cagc	agca	gccaca	420
ttt	cagt	tac	agct	ccas	gac t	ccca	ggtt	g ca	agact	ccca	aag	gcaaa	açag	acto	cagtcc	480
acç	gated	agc	tcca	agtto	ca c	cgat	ccga	at co	cacto	gatac	ago	gtg	ctcg	agtg	gccatag	540
ato	ctca	acca	agt	gcicaa	aa t	ccg	atco	et ga	atcco	caaga	gct	caa	ggca	ccc	ggccca	600

aaattgagct gagaacgaaa cgaaggaagt tccttagtgc catagaaagc agttaatgaa

660

acaacgacta agacgaagat cgaccatcca gaaccggagg gagctaattg cgaacgaaag 720 aaaccacaaa gtgccttcca tcaatccgtt gataagtgat atttattatg tttatacttg 780 ccagcagccg aggcagcaac agcaatagca acaaccatag gggatcacgg catcgatgat 840 cagtccacga ccaagtccta gtgcaatccg gaatccagtt caaattagtt caa'taagccg 900 tatctaccac gtataatgtc cacatccacc gccacaacga gcgttatcac gtccaacgag 960 etetegetgt ceggecaege ceaeggteae ggteaegeee accagttgea ceageaeaee 1020 cacagoogco taggagttgg cgttggtgtt ggcatootta gcgacgcato cotatogcoc 1080 atccaacaag gcagtggcgg ccacagcggc ggaggtaaca caaacagttc accactggcg 1140 cccaacggag tgccacttct cacaacaatg caccgatcac cggactcacc gcagccagaa 1200 ttggccacca tgacgaacgt caacgtgctg gatctgcaca cggataactc caagctgtac 1260 gacaaggagg ctgtatttat atacgaaacg cccaaggtgg tgatgccagc ggatggcggg 1320 ggtggcaata attccgatga aggtcatgcc atcgatgcgc ggattgcggc ccaaatgggc 1380 aaccaagece ageaacagea geageageaa cageagaegg aacaceagee getggeeaag 1440 atcgagttcg atgagaacca gataatccgg gtggtgggac caaatggcga gcaacagcaa 1500 atcatctcgc gggagatcat caatggggag catcatatcc tgtcgcgaaa cgaggctggt 1560 gagcacattc tcacacggat cgtcagtgat ccctccaagt tgatgcccaa tgacaatgca 1620 gtggccacgg ccatgtacaa ccaggcccaa aagatgaaca atgatcacgg gcaggcggta 1680 tatcagacat caccattgcc gctagacgcg tctgtattgc attatagtgg cggcaatgat 1740 tegaatgtaa ttaagaegga ggeegatate taegaggate acaagaaaca tgeggetgea 1800 gcagcagctg ctgccggcgg aggatccatc atatacacca catccgatcc gaacggagtg 1860 aatgtgaaac aactgcccca tttgacggta ccccaaaaac ttgatcccga cctctatcaa 1920 gccgataagc atatagattt gatctacaac gatggcagca agacggtgat ttactccact 1980 acggatcaga agagtttgga aatatactcg ggcggcgaca tcggcagcct ggtgtccgac 2040 ggccaagtgg tggtccaggc gggactgccg tatgccacca ccaccggagc cggcggccag 2100 cccgtctata tcgtggccga cggtgccttg ccagcgggag tcgaggagca tctgcagagt 2160 ggaaagetea atggecagae cacacetate gatgtetetg geetategea aaatgagatt 2220 caaggetttt tgeteggete acaeceeteg teateggega eggtaageac aaceggegtt 2280 gtctccacga caacgatctc gcatcaccag caacagcagc agcagcagca acagcaacag 2340

cagcagcagc	agcagcaaca	ccagcagcag	cagcaacatc	ccggcgacat	tgttagtgcc	2400
gctggcgtgg	ggagcacggg	ctccattgtc	tectetgegg	cgcaacagca	gcagcagcag	2460
caactaatta	gcatcaaacg	agagcccgaa	gacttgcgca	aggatcccaa	gaatggcaac	2520
attgccggtg	cagcaacagc	aaatggaccc	ggttcggtca	taacccaaaa	gatcttgcac	2580
gtggatgcac	caacggcaag	tgaagctgat	aggcccagca	cacccagcag	cagcatcaac	2640
agcactgaaa	acactgaatc	ggactcacag	tcagtatcag	gatcagaatc	aggatcgccg	2700
ggagccagga	ccacagccac	actagagatg	tatgcaacca	cgggcggcac	acagatctat	2760
ctacagacct	cacatcccag	cacggcgagc	ggagcgggcg	acaacaccaa	accegetgga	2820
gccgccggcg	gcggcggtgt	gtccatgcag	gcgcaaagtc	ccagtccggg	tccctatatc	2880
acggccaatg	actatggcat	gtacacggcc	agtcgcctgc	cacccggtcc	cccgcccacc	2940
agcaccacca	cgtttatagc	ggagccctcc	tactatcggg	aatactttgc	accggatggc	3000
caaggtggct	atgtgccggc	cagcacgagg	totttgtatg	gcgacgtgga	cgtatccgta	3060
tctcagcccg	gcggagtggt	cacctatgag	ggccgctttg	ccggcagcgt	tccccgccc	3120
gccaccacca	ccgtgctaac	cagcgtgcat	caccaccagc	aacagcagca	gcaacaacag	3180
cagcatcaac	agcagcagca	gcagcaacag	caccaccagc	agcaacagca	ccattcgcag	3240
gatggcaaga	gcaatggcgg	agcaacgcca	ctctatgcca	aagccattac	ggcggcgggt	3300
ctaacggtgg	g atttgccaag	teeggatteg	ggcattggta	cggatgccat	tacaccgcgg	3360
gatcagacaa	atatccaaca	gtcctttgat	tatacggaat	tgtgccagcc	gggcacgctg	3420
atcgatgcca	a atggcagcat	accegtcage	gtgaacagca	tccagcagag	aacggcggtc	3480
catggcagco	e agaacagtco	caccacatcg	ctggtggaca	ccagcaccaa	tggatccacg	3540
cgatcgcgg	c cctggcacga	ctttggacgt	: cagaatgatg	ccgacaaaat	acaaatacca	3600
aaaatcttc	a caaacgtggg	cttccgatat	. cacctggaga	gccccatcag	ttcatcgcag	3660
aggcgcgag	g acgatcgcat	cacctacato	: aacaagggto	: aattctatgg	aataacgctg	3720
gagtatgtg	c acgatgcgga	aaagcccatt	: aagaacacca	ccgtcaagag	, tgtgatcatg	3780
ctaatgttc	c gcgaggagaa	a gagtcccgag	g gatgagatca	aggeetggea	attctggcac	3840
agtcgtcag	c attccgtga	a gcagagaato	ttggatgcag	g atacgaagaa	e ctcggttggc	3900
ctcgttggc	t gcatcgagga	a agtgtcgca	c aatgccatco	g ccgtctactg	g gaatccgctg	3960
gagagetee	g ccaagatca	a cattgcggt	t cagtgettga	a gcacggattt	cagcagtcaa	4020

aagggaggcc tgccgctgca cgtacaaatc gacacatttg aggaccccag agatacggcg 4080 gtcttccacc gcggctactg tcagataaag gtcttctgcg ataagggcgc cgaacgaaag 4140 acgegegatg aagageggeg ggeegeeaaa egaaagatga cageeaeggg cagaaagaag 4200 ctggacgage tttaccatcc ggtaacggat cggtccgagt tctatggcat gcaggacttc 4260 gccaagccgc cggtgctatt ctcgcccgcc gaggacatgg agaaggtagg tcagctgggc 4320 attggcgctg ccaccggcat gacattcaac cccctgagca acggcaactc caactccaac 4380 tegeactegt cettgeagag ettetaegge catgagaetg actegeegga eetgaagggg 4440 gcctcaccgt tcctgctcca cggccagaag gtggccacgc cgacgctcaa gttccacaac 4500 cattttccgc ccgacatgca gaccgataag aaggatcaca tactggacca gaacatgttg 4560 accagcacac ccctgaccga ctttggtccg ccgatgaagc gcggcaggat gacgccgccg 4620 accteggaae gegtgatget gtaegtgegg eaggagaaeg aggaggtgta tacacegttg 4680 cacgtggtgc cgcccacçac gatcggcctg ctaaatgcga ttgaaaacaa atacaaaatc 4740 tcaacaacga gcataaataa catttatcgc acaaacaaga aggggattac tgcgaaaatt 4800 gacgatgaca tgatatcgtt ctactgcaac gaggacatct ttctgctgga ggtgcaacag 4860 atcgaggacg acctgtacga tgtgacgete acggagetge ceaatcagta gegetggeag 4920 tacgggtagc acccgctaac cgcactcaaa aaaaaaagca aacaaacaca caaattacgg 4980 acacaacaag ttgtttcaat aagccatttt ccatagagcc taagtctaaa tatcgtagtt 5040 ataataatgg gatccgcaac aaatcgagtt gcaacgaatg ttaagaacgc taacacaata 5100 cgcatgtaaa atgatacttt aaaattgatt tagttatttt agcaacaatg agattatcta 5160 aaattgtttg atcaaatttt acattctcgc tatgtctata gataattcta agcccgtaag 5220 cccataagcg taatcgtaat cgtaatcgta ccgtgtattt atgctcatat ataaacaact 5280 atatatata atatatat atatatgtgc ggagtgcaac agtgtctgtc cagtaggaga 5340 taagtotogt ttoogotoco otgottatgo tatgacotta ggtocagggo aagtatgagt 5400 taccgaatct atctattagg tgcatctaac gaaaggaatc attagctctg cacgaactct 5460 agccgtagcc tattgtaatc catttgtatg tttggcttaa gcgttttact tgttgaatat 5520 aaagtgtaaa attatttttg aaaaaaaaaa acccacacaa aacacaaatc gtttgttcta 5580 tatttctgtt tcaaaactaa ctcgttaccc acaatcccct ctgttatgta taattaggat 5640 ctctgtacac 5650

-

<210> 37

<211> 1331 <212> PRT <213> Drosophila <400> Met Ser Thr Ser Thr Ala Thr Thr Ser Val Ile Thr Ser Asn Glu Leu Ser Leu Ser Gly His Ala His Gly His Gly His Ala His Gln Leu His Gln His Thr His Ser Arg Leu Gly Val Gly Val Gly Val Gly Ile Leu Ser Asp Ala Ser Leu Ser Pro Ile Gln Gln Gly Ser Gly Gly His Ser Gly Gly Asn Thr Asn Ser Ser Pro Leu Ala Pro Asn Gly Val Pro Leu Leu Thr Thr Met His Arg Ser Pro Asp Ser Pro Gln Pro Glu Leu 90 Ala Thr Met Thr Asn Val Asn Val Leu Asp Leu His Thr Asp Asn Ser Lys Leu Tyr Asp Lys Glu Ala Val Phe Ile Tyr Glu Thr Pro Lys Val 120 Val Met Pro Ala Asp Gly Gly Gly Asn Asn Ser Asp Glu Gly His Ala Ile Asp Ala Arg Ile Ala Ala Gln Met Gly Asn Gln Ala Gln Gln Gln Gln Gln Gln Gln Gln Thr Glu His Gln Pro Leu Ala Lys Ile Glu Phe Asp Glu Asn Gln Ile Ile Arg Val Val Gly Pro Asn Gly Glu Gln Gln Gln Ile Ile Ser Arg Glu Ile Ile Asn Gly Glu His His Ile 195 200 Leu Ser Arg Asn Glu Ala Gly Glu His Ile Leu Thr Arg Ile Val Ser Asp Pro Ser Lys Leu Met Pro Asn Asp Asn Ala Val Ala Thr Ala Met Tyr Asn Gln Ala Gln Lys Met Asn Asn Asp His Gly Gln Ala Val Tyr 245 250 Gln Thr Ser Pro Leu Pro Leu Asp Ala Ser Val Leu His Tyr Ser Gly 270 265

Gly Asn Asp Ser Asn Val Ile Lys Thr Glu Ala Asp Ile Tyr Glu Asp His Lys Lys His Ala Ala Ala Ala Ala Ala Ala Gly Gly Gly Ser Ile Ile Tyr Thr Thr Ser Asp Pro Asn Gly Val Asn Val Lys Gln Leu 310 Pro His Leu Thr Val Pro Gln Lys Leu Asp Pro Asp Leu Tyr Gln Ala 330 Asp Lys His Ile Asp Leu Ile Tyr Asn Asp Gly Ser Lys Thr Val Ile 345 Tyr Ser Thr Thr Asp Gln Lys Ser Leu Glu Ile Tyr Ser Gly Gly Asp 355 Ile Gly Ser Leu Val Ser Asp Gly Gln Val Val Val Gln Ala Gly Leu Pro Tyr Ala Thr Thr Gly Ala Gly Gly Gln Pro Val Tyr Ile Val Ala Asp Gly Ala Leu Pro Ala Gly Val Glu Glu His Leu Gln Ser Gly Lys Leu Asn Gly Gln Thr Thr Pro Ile Asp Val Ser Gly Leu Ser Gln 425 Asn Glu Ile Gln Gly Phe Leu Leu Gly Ser His Pro Ser Ser Ser Ala Thr Val Ser Thr Thr Gly Val Val Ser Thr Thr Thr Ile Ser His His 465 Gln His Gln Gln Gln Gln His Pro Gly Asp Ile Val Ser Ala Ala 490 Gly Val Gly Ser Thr Gly Ser Ile Val Ser Ser Ala Ala Gln Gln 500 Gln Gln Gln Leu Ile Ser Ile Lys Arg Glu Pro Glu Asp Leu Arg 520 Lys Asp Pro Lys Asn Gly Asn Ile Ala Gly Ala Ala Thr Ala Asn Gly 535 Pro Gly Ser Val Ile Thr Gln Lys Ile Leu His Val Asp Ala Pro Thr Ala Ser Glu Ala Asp Arg Pro Ser Thr Pro Ser Ser Ser Ile Asn Ser

575 565 570 Thr Glu Asn Thr Glu Ser Asp Ser Gln Ser Val Ser Gly Ser Glu Ser 585 Gly Ser Pro Gly Ala Arg Thr Thr Ala Thr Leu Glu Met Tyr Ala Thr Thr Gly Gly Thr Gln Ile Tyr Leu Gln Thr Ser His Pro Ser Thr Ala Ser Gly Ala Gly Gly Gly Ala Gly Pro Ala Gly Ala Ala Gly Gly Gly 630 625 Gly Val Ser Met Gln Ala Gln Ser Pro Ser Pro Gly Pro Tyr Ile Thr 650 Ala Asn Asp Tyr Gly Met Tyr Thr Ala Ser Arg Leu Pro Pro Gly Pro Pro Pro Thr Ser Thr Thr Thr Phe Ile Ala Glu Pro Ser Tyr Tyr Arg 680 Glu Tyr Phe Ala Pro Asp Gly Gln Gly Gly Tyr Val Pro Ala Ser Thr Arg Ser Leu Tyr Gly Asp Val Asp Val Ser Val Ser Gln Pro Gly Gly Val Val Thr Tyr Glu Gly Arg Phe Ala Gly Ser Val Pro Pro Ala 730 Thr Thr Thr Val Leu Thr Ser Val His His His Gln Gln Gln Gln 745 Gln Gln Gln His Gln Gln Gln Gln Gln Gln Gln His His Gln 760 Gln Gln Gln His His Ser Gln Asp Gly Lys Ser Asn Gly Gly Ala Thr Pro Leu Tyr Ala Lys Ala Ile Thr Ala Ala Gly Leu Thr Val Asp Leu Pro Ser Pro Asp Ser Gly Ile Gly Thr Asp Ala Ile Thr Pro Arg Asp 805 810 Gln Thr Asn Ile Gln Gln Ser Phe Asp Tyr Thr Glu Leu Cys Gln Pro Gly Thr Leu Ile Asp Ala Asn Gly Ser Ile Pro Val Ser Val Asn Ser 840 Ile Gln Gln Arg Thr Ala Val His Gly Ser Gln Asn Ser Pro Thr Thr 855

Ser Leu Val Asp Thr Ser Thr Asn Gly Ser Thr Arg Ser Arg Pro Trp

His Asp Phe Gly Arg Gln Asn Asp Ala Asp Lys Ile Gln Ile Pro Lys 885 Ile Phe Thr Asn Val Gly Phe Arg Tyr His Leu Glu Ser Pro Ile Ser Ser Ser Gln Arg Arg Glu Asp Asp Arg Ile Thr Tyr Ile Asn Lys Gly Gln Phe Tyr Gly Ile Thr Leu Glu Tyr Val His Asp Ala Glu Lys Pro Ile Lys Asn Thr Thr Val Lys Ser Val Ile Met Leu Met Phe Arg Glu Glu Lys Ser Pro Glu Asp Glu Ile Lys Ala Trp Gln Phe Trp His Ser 965 Arg Gln His Ser Val Lys Gln Arg Ile Leu Asp Ala Asp Thr Lys Asn Ser Val Gly Leu Val Gly Cys Ile Glu Glu Val Ser His Asn Ala Ile 995 Ala Val Tyr Trp Asn Pro Leu Glu Ser Ser Ala Lys Ile Asn Ile 1015 1020 1010 Ala Val Gln Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Gly 1030 Leu Pro Leu His Val Gln Ile Asp Thr Phe Glu Asp Pro Arg Asp Thr Ala Val Phe His Arg Gly Tyr Cys Gln Ile Lys Val Phe Cys 1065 1055 1060 Asp Lys Gly Ala Glu Arg Lys Thr Arg Asp Glu Glu Arg Arg Ala 1070 Ala Lys Arg Lys Met Thr Ala Thr Gly Arg Lys Lys Leu Asp Glu Leu Tyr His Pro Val Thr Asp Arg Ser Glu Phe Tyr Gly Met Gln 1100 1105 1110 Asp Phe Ala Lys Pro Pro Val Leu Phe Ser Pro Ala Glu Asp Met 1120 Glu Lys Val Gly Gln Leu Gly Ile Gly Ala Ala Thr Gly Met Thr 1135 Phe Asn Pro Leu Ser Asn Gly Asn Ser Asn Ser Asn Ser His Ser 1145 1150

Ser	Leu 1160	Gln	Ser	Phe	Tyr	Gly 1165	His	Glu	Thr	Asp	Ser 1170	Pro	Asp	Leu	
Lys	Gly 1175		Ser	Pro	Phe	Leu 1180		His	Gly	Gln	Lys 1185	Val	Ala	Thr	
Pro	Thr 1190		Lys	Phe	His	Asn 1195		Phe	Pro	Pro	Asp 1200	Met	Gln	Thr	
Asp	Lys 1205	_	Asp	His	Ile	Leu 1210		Gln	Asn	Met	Leu 1215	Thr	Ser	Thr	
Pro	Leu 1220		Asp	Phe	Gly	Pro 1225		Met	Lys	Arg	Gly 1230	Arig	Met	Thr	
Pro	Pro 1235		Ser	Glu	Arg	Val 1240		Leu	тут	Vạl	Arg 1245		Glu	Asn	
Glu	Glu 1250		Tyr	Thr	Pro	Leu 1255		Val	Val	Pro	Pro 1260		Thr	Ile	
Gly	Leu 1265		Asn	Ala	Ile	Glu 1270		Lys	Туг	. Lys	Ile 1275	Ser	Thr	Thr	
Ser	Ile 1280		Asn	Ile	Tyr	Arg 1285		Asn	Lys	r PÀa	Gly 1290	Ile	Thr	Ala	
Lys	Ile 1295		Asp	Asp	Met	Ile 1300		Phe	туг	Cys	1305	Glu	. Asp	) Ile	
Phe	Leu 1310		Glu	. Val	Gln	Gln 1315		e Glu	ı Ası	) Asp	1320	тут	Asp	Val	
Thr	Leu 1325		Glu	ı Lev	Pro	) Asn 1330		1							
<21	LO> 3	38													
	L1> 5														
	L2> I L3> I		ophil	la				•	•	•					
			-									•			
aaa	00> : aaata	gaa a	aaaa	caaca	aa c	aaatt	ggct	tga	aaac	gca	aatgc	cagg	g gc	aacgcccc	6
cga	aaccg	acc (	cgcc	ccct	ca a	etttt	gcgc	cct	ccag	tag	caata	gcag	c aa	tatgagca	12
gc	agcaa	cat (	caaa	tgtt	ag g	ccaaa	atgc	aca	aacc	gcc ·	agcaa	caaa	g gc	agcaccaa	18
gc	gaacg	aaa	caac	aaca	gc t	ccaca	tacc	aca	aaga	gtg	gcaca	ttag	a ag	cggccaaa	24
ag	cagcc	agc	cgag	agca	tt g	tgtaa	gcca	aag	gccc	aga	gagcc	aggc	t aa	aagccccc	30
ag	acgca	caa	caac	aaca	ac a	acaac	taaa	aca	.gcac	aaa	gagtg	gcga	a ag	gtgcaccc	36
ac	cagca	aaa	cagc	aaca	ac g	gagca	.acca	aca	acag	cag	cagca	gcag	c ag	cagccaca	42



cccgtctata	tegtggeega	eggtgeettg	ccagcgggag	tcgaggagca	tctgcagagt	2160
ggaaagctca	atggccagac	cacacctatc	gatgtctctg	gcctatcgca	aaatgagatt	2220
caaggetttt	tgctcggctc	acacccctcg	tcatcggcga	cggtaagcac	aaccggcgtt	2280
gtctccacga	caacgatctc	gcatcaccag	caacagcagc	agcagcagca	acagcaacag	2340
cagcagcagc	agcagcaaca	ccagcagcag	cagcaacatc	ccggcgacat	tgttagtgcc	2400
gctggcgtgg	ggagcacggg	ctccattgtc	tectetgegg	cgcaacagca	gcagcagcag	2460
caactaatta	gcatcaaacg	agagcccgaa	gacttgcgca	aggatcccaa	gaatggcaac	2520
attgccggtg	cagcaacagc	aaatggaccc	ggttcggtca	taacccaaaa	gatcttgcac	2580
gtggatgcac	caacggcaag	tgaagctgat	aggcccagca	cacccagcag	cagcatcaac	2640
agcactgaaa	acactgaatc	ggactcacag	tcagtatcag	gatcagaatc	aggatcgccg	2700
ggagçcagga	ccacagccac	actagagatg	tatgcaacca	cgggcggcac	acagatetat	2760
ctacagacct	cacatcccag	cacggcgagc	ggagcgggcg	geggegeegg	acccgctgga	2820
gccgccggcg	geggeggtgt	gtccatgcag	gcgcaaagtc	ccagtccggg	tccctatatc	2880
acggccaatg	actatggcat	gtacacggcc	agtcgcctgc	cacccggtcc	cccgcccacc	2940
agcaccacca	cgtttatago	ggagccctcc	tactatcggg	aatactttgc	accggatggc	3000
caaggtggct	: atgtgccggc	cagcacgagg	tctttgtatg	gcgacgtgga	cgtatccgta	3060
teteageeeg	g gcggagtggt	cacctatgag	ggccgctttg	ccggcagcgt	teeceegeee	3120
gccaccacca	ccgtgctaac	: cagcgtgcat	caccaccagc	aacagcagca	gcaacaacag	3180
cagcatcaac	agcagcagca	gcagcaacag	r çaccaccago	agcaacagca	ccattcgcag	3240
gatggcaaga	a gcaatggcgg	g agcaacgcca	ctctatgcca	aagccattac	: ggcggcgggt	3300
ctaacggtg	g atttgccaag	g teeggatteg	g ggcattggta	cggatgccat	: tacaccgcgg	3360
gatcagaca	a atatccaaca	a gtcctttgat	: tatacggaat	: tgtgccagco	gggcacgctg	3420
atcgatgcc	a atggcagca	t acceptcage	gtgaacagca	tccagcagag	g aacggcggtc	3480
catggcagc	c agaacagtc	c caccacatco	g ctggtggaca	a ccagcacca	a tggatccacg	3540
cgatcgcgg	c cctggcacg	a ctttggacgi	t cagaatgatg	g ccgacaaaa	t acaaatacca	3600
aaaatcttc	a caaacgtgg	g cttccgata	t cacctggaga	a gccccatca	g ttcatcgcag	3660
aggcgcgag	g acgatcgca	t cacctacat	c aacaagggt	c aattotatg	g aataacgctg	3720
gagtatgtg	c acgatgcgg	a aaagcccat	t aagaacacca	a ccgtcaaga	g tgtgatcatg	3780




cacacaaaac acaaatcgtt tgttctatat ttctgtttca aaactaactc gttacccaca 5520 atcccctctg ttatgtataa ttaggatctc tgtacac 5557

<210> 39 <211> 1331 PRT <213> Drosophila <400> 39 Met Ser Thr Ser Thr Ala Thr Thr Ser Val Ile Thr Ser Asn Glu Leu Ser Leu Ser Gly His Ala His Gly His Gly His Ala His Gln Leu His Gln His Thr His Ser Arg Leu Gly Val Gly Val Gly Val Gly Ile Leu Ser Asp Ala Ser Leu Ser Pro Ile Gln Gly Ser Gly Gly His Ser Gly Gly Asn Thr Asn Ser Ser Pro Leu Ala Pro Asn Gly Val Pro Leu Leu Thr Thr Met His Arg Ser Pro Asp Ser Pro Gln Pro Glu Leu 90 Ala Thr Met Thr Asn Val Asn Val Leu Asp Leu His Thr Asp Asn Ser Lys Leu Tyr Asp Lys Glu Ala Val Phe Ile Tyr Glu Thr Pro Lys Val 120 Val Met Pro Ala Asp Gly Gly Gly Asn Asn Ser Asp Glu Gly His Ala Ile Asp Ala Arg Ile Ala Ala Gln Met Gly Asn Gln Ala Gln Gln Gln Gln Gln Gln Gln Gln Thr Glu His Gln Pro Leu Ala Lys Ile 170 Glu Phe Asp Glu Asn Gln Ile Ile Arg Val Val Gly Pro Asn Gly Glu Gln Gln Gln Ile Ile Ser Arg Glu Ile Ile Asn Gly Glu His His Ile Leu Ser Arg Asn Glu Ala Gly Glu His Ile Leu Thr Arg Ile Val Ser 215 Asp Pro Ser Lys Leu Met Pro Asn Asp Asn Ala Val Ala Thr Ala Met 235

Tyr Asn Gln Ala Gln Lys Met Asn Asn Asp His Gly Gln Ala Val Tyr

250 Gln Thr Ser Pro Leu Pro Leu Asp Ala Ser Val Leu His Tyr Ser Gly Gly Asn Asp Ser Asn Val Ile Lys Thr Glu Ala Asp Ile Tyr Glu Asp 280 His Lys Lys His Ala Ala Ala Ala Ala Ala Ala Gly Gly Gly Ser Ile Ile Tyr Thr Thr Ser Asp Pro Asn Gly Val Asn Val Lys Gln Leu 310 Pro His Leu Thr Val Pro Gln Lys Leu Asp Pro Asp Leu Tyr Gln Ala 330 Asp Lys His Ile Asp Leu Ile Tyr Asn Asp Gly Ser Lys Thr Val Ile Tyr Ser Thr Thr Asp Gln Lys Ser Leu Glu Ile Tyr Ser Gly Gly Asp 355 Ile Gly Ser Leu Val Ser Asp Gly Gln Val Val Gln Ala Gly Leu Pro Tyr Ala Thr Thr Gly Ala Gly Gly Gln Pro Val Tyr Ile Val Ala Asp Gly Ala Leu Pro Ala Gly Val Glu Glu His Leu Gln Ser Gly 410 405 Lys Leu Asn Gly Gln Thr Thr Pro Ile Asp Val Ser Gly Leu Ser Gln Asn Glu Ile Gln Gly Phe Leu Leu Gly Ser His Pro Ser Ser Ser Ala Thr Val Ser Thr Thr Gly Val Val Ser Thr Thr Thr Ile Ser His His Gln His Gln Gln Gln Gln His Pro Gly Asp Ile Val Ser Ala Ala 490 Gly Val Gly Ser Thr Gly Ser Ile Val Ser Ser Ala Ala Gln Gln 500 Gln Gln Gln Leu Ile Ser Ile Lys Arg Glu Pro Glu Asp Leu Arg Lys Asp Pro Lys Asn Gly Asn Ile Ala Gly Ala Ala Thr Ala Asn Gly

535 540 530 Pro Gly Ser Val Ile Thr Gln Lys Ile Leu His Val Asp Ala Pro Thr Ala Ser Glu Ala Asp Arg Pro Ser Thr Pro Ser Ser Ser Ile Asn Ser 565 570 Thr Glu Asn Thr Glu Ser Asp Ser Gln Ser Val Ser Gly Ser Glu Ser Gly Ser Pro Gly Ala Arg Thr Thr Ala Thr Leu Glu Met Tyr Ala Thr 600 Thr Gly Gly Thr Gln Ile Tyr Leu Gln Thr Ser His Pro Ser Thr Ala Ser Gly Ala Gly Gly Gly Ala Gly Pro Ala Gly Ala Ala Gly Gly Gly Gly Val Ser Met Gln Ala Gln Ser Pro Ser Pro Gly Pro Tyr Ile Thr 650 Ala Asn Asp Tyr Gly Met Tyr Thr Ala Ser Arg Leu Pro Pro Gly Pro Pro Pro Thr Ser Thr Thr Thr Phe Ile Ala Glu Pro Ser Tyr Tyr Arg Glu Tyr Phe Ala Pro Asp Gly Gln Gly Gly Tyr Val Pro Ala Ser Thr 695 Arg Ser Leu Tyr Gly Asp Val Asp Val Ser Val Ser Gln Pro Gly Gly Val Val Thr Tyr Glu Gly Arg Phe Ala Gly Ser Val Pro Pro Pro Ala Thr Thr Thr Val Leu Thr Ser Val His His His Gln Gln Gln Gln 745 Gln Gln Gln His Gln Gln Gln Gln Gln Gln Gln His His Gln Gln Gln Gln His His Ser Gln Asp Gly Lys Ser Asn Gly Gly Ala Thr Pro Leu Tyr Ala Lys Ala Ile Thr Ala Ala Gly Leu Thr Val Asp Leu Pro Ser Pro Asp Ser Gly Ile Gly Thr Asp Ala Ile Thr Pro Arg Asp 810 Gln Thr Asn Ile Gln Gln Ser Phe Asp Tyr Thr Glu Leu Cys Gln Pro



- Ile Gln Gln Arg Thr Ala Val His Gly Ser Gln Asn Ser Pro Thr Thr 850 855 860
- Ser Leu Val Asp Thr Ser Thr Asn Gly Ser Thr Arg Ser Arg Pro Trp 865 870 875 880
- His Asp Phe Gly Arg Gln Asn Asp Ala Asp Lys Ile Gln Ile Pro Lys 885 890 895
- Ile Phe Thr Asn Val Gly Phe Arg Tyr His Leu Glu Ser Pro Ile Ser 900 905 910
- Ser Ser Gln Arg Arg Glu Asp Asp Arg Ile Thr Tyr Ile Asn Lys Gly 915 920 925
- Gln Phe Tyr Gly Ile Thr Leu Glu Tyr Val His Asp Ala Glu Lys Pro 930 935 940
- Ile Lys Asn Thr Thr Val Lys Ser Val Ile Met Leu Met Phe Arg Glu 945 950 955 960
- Glu Lys Ser Pro Glu Asp Glu Ile Lys Ala Trp Gln Phe Trp His Ser 965 970 975
- Arg Gln His Ser Val Lys Gln Arg Ile Leu Asp Ala Asp Thr Lys Asn 980 985 990
- Ser Val Gly Leu Val Gly Cys Ile Glu Glu Val Ser His Asn Ala Ile 995 1000 1005
- Ala Val Tyr Trp Asn Pro Leu Glu Ser Ser Ala Lys Ile Asn Ile 1010 1015 1020
- Ala Val Gln Cys Leu Ser Thr Asp Phe Ser Ser Gln Lys Gly Gly 1025 1030 1035
- Leu Pro Leu His Val Gln Ile Asp Thr Phe Glu Asp Pro Arg Asp 1040 1045 1050
- Thr Ala Val Phe His Arg Gly Tyr Cys Gln Ile Lys Val Phe Cys 1055 1060 1065
- Asp Lys Gly Ala Glu Arg Lys Thr Arg Asp Glu Glu Arg Arg Ala

1070 1075 Ala Lys Arg Lys Met Thr Ala Thr Gly Arg Lys Leu Asp Glu Leu Tyr His Pro Val Thr Asp Arg Ser Glu Phe Tyr Gly Met Gln - 1110 Asp Phe Ala Lys Pro Pro Val Leu Phe Ser Pro Ala Glu Asp Met Glu Lys Val Gly Gln Leu Gly Ile Gly Ala Ala Thr Gly Met Thr Phe Asn Pro Leu Ser Asn Gly Asn Ser Asn Ser Asn Ser His Ser Ser Leu Gln Ser Phe Tyr Gly His Glu Thr Asp Ser Pro Asp Leu Lys Gly Ala Ser Pro Phe Leu Leu His Gly Gln Lys Val Ala Thr Pro Thr Leu Lys Phe His Asn His Phe Pro Pro Asp Met Gln Thr Asp Lys Lys Asp His Ile Leu Asp Gln Asn Met Leu Thr Ser Thr Pro Leu Thr Asp Phe Gly Pro Pro Met Lys Arg Gly Arg Met Thr Pro Pro Thr Ser Glu Arg Val Met Leu Tyr Val Arg Gln Glu Asn Glu Glu Val Tyr Thr Pro Leu His Val Val Pro Pro Thr Thr Ile Gly Leu Leu Asn Ala Ile Glu Asn Lys Tyr Lys Ile Ser Thr Thr Ser Ile Asn Asn Ile Tyr Arg Thr Asn Lys Lys Gly Ile Thr Ala - 74 -

1280 1285 1290

Lys Ile Asp Asp Asp Met Ile Ser Phe Tyr Cys Asn Glu Asp Ile 1295 1300 1305

Phe Leu Leu Glu Val Gln Gln Ile Glu Asp Asp Leu Tyr Asp Val 1310 1315 1320

Thr Leu Thr Glu Leu Pro Asn Gln 1325 1330

<u>DATED</u> this twenty-second day of August 2002.

## Melbourne Health

by DAVIES COLLISION CAVE

Patent Attorneys for the Applicant

## This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
□ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
□ FADED TEXT OR DRAWING
□ BLURRED OR ILLEGIBLE TEXT OR DRAWING
□ SKEWED/SLANTED IMAGES
□ COLOR OR BLACK AND WHITE PHOTOGRAPHS
□ GRAY SCALE DOCUMENTS
□ LINES OR MARKS ON ORIGINAL DOCUMENT
□ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

## IMAGES ARE BEST AVAILABLE COPY.

☐ OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.